

## On some methods of probability

Alžbeta Michalíková<sup>1</sup> and Beloslav Riečan<sup>1,2</sup>

<sup>1</sup> Faculty of Natural Sciences, Matej Bel University  
Tajovského 40, Banská Bystrica, Slovakia  
e-mail: alzbeta.michalikova@umb.sk

<sup>2</sup> Mathematical Institute, Slovak Academy of Sciences  
Ďumbierska 1, Banská Bystrica, Slovakia  
e-mail: beloslav.riecan@umb.sk

**Received:** 28 March 2018

**Accepted:** 20 April 2018

**Abstract:** The paper contains a review of some methods for building probability theory on intuitionistic fuzzy sets. They are based on some representation of states by Kolmogorov probability spaces as well as the embedding of  $IF$ -spaces into the  $MV$ -algebras.

**Keywords:**  $IF$ -sets,  $IF$ -states,  $IV$ -sets,  $MV$ -algebras.

**2010 Mathematics Subject Classification:** 03E72.

### 1 Introduction

In this paper we will work with probability defined on intuitionistic fuzzy sets (shortly  $IF$ -sets). First we will describe the structures which we will work with. Then the relations between the different structures and their properties will be showed. Let start with the definition of the important term in probability theory - state. Consider a measurable space  $(\Omega, \mathcal{S})$  with a  $\sigma$ -algebra  $\mathcal{S}$ . Let  $\mathcal{J}$  be the family of all measurable functions  $f : \Omega \rightarrow [0, 1]$ . To define the state on  $\mathcal{J}$  we need two binary operations. In this paper we will use the Lukasiewicz operations

$$f \oplus g = \min(f + g, 1_\Omega),$$

$$f \odot g = \max(f + g - 1_\Omega, 0_\Omega).$$

These binary operations play the same role as the union and the intersection in the set theory.

A state on  $\mathcal{J}$  is a mapping  $m_{\mathcal{J}} : \mathcal{J} \rightarrow [0, 1]$  satisfying the following conditions

1.  $m_{\mathcal{J}}(0_{\Omega}) = 0, m_{\mathcal{J}}(1_{\Omega}) = 1.$
2. If  $f \odot g = 0_{\Omega}$  then  $m_{\mathcal{J}}(f \oplus g) = m_{\mathcal{J}}(f) + m_{\mathcal{J}}(g).$
3. If  $f_n \nearrow f$  then  $m_{\mathcal{J}}(f_n) \nearrow m_{\mathcal{J}}(f).$

Since  $\mathcal{J}$  contains the functions  $f : \Omega \rightarrow [0, 1]$  then  $\mathcal{J}$  represent the family of fuzzy sets. In the fuzzy set theory  $f$  is called membership function. Let us now look at the more general structure - intuitionistic fuzzy sets. The Atanassov *IF*-set [1, 2] is a pair  $A = (\mu_A, \nu_A)$  of fuzzy sets  $\mu_A : \Omega \rightarrow [0, 1], \nu_A : \Omega \rightarrow [0, 1]$  such that  $\mu_A + \nu_A \leq 1_{\Omega}$ . Similarly as in the fuzzy set theory the function  $\mu_A$  is called the membership function. The second function,  $\nu_A$  is called the non-membership function. Denote by  $\mathcal{F}$  the family of all *IF*-sets such that  $\mu_A, \nu_A$  are  $\mathcal{S}$ -measurable. On the set  $\mathcal{F}$  there is defined the following ordering

$$A = (\mu_A, \nu_A) \leq (\mu_B, \nu_B) = B \iff \mu_A \leq \mu_B, \nu_A \geq \nu_B.$$

By this ordering for each  $A \in \mathcal{F}$  it holds

$$(0_{\Omega}, 1_{\Omega}) \leq A \leq (1_{\Omega}, 0_{\Omega}).$$

The family  $\mathcal{J}$  can be considered as a subset of  $\mathcal{F}$  if for  $f \in \mathcal{J}$  we put  $A = (f, 1 - f)$ . Then  $\mu_A + \nu_A = 1$ .

For  $A, B \in \mathcal{F}$  we could define the Lukasiewicz operations

$$A \oplus B = (\min(\mu_A + \mu_B, 1_{\Omega}), \max(\nu_A + \nu_B - 1_{\Omega}, 0_{\Omega})),$$

$$A \odot B = (\max(\mu_A + \mu_B - 1_{\Omega}, 0_{\Omega}), \min(\nu_A + \nu_B, 1_{\Omega})).$$

A state on  $\mathcal{F}$  is a mapping  $m : \mathcal{F} \rightarrow [0, 1]$  satisfying the following conditions [6]

1.  $m((0_{\Omega}, 1_{\Omega})) = 0, m((1_{\Omega}, 0_{\Omega})) = 1,$
2.  $A \odot B = (0_{\Omega}, 1_{\Omega}) \implies m(A \oplus B) = m(A) + m(B),$
3.  $A_n \nearrow A \implies m(A_n) \nearrow m(A).$

In this contribution we will also work with *MV*-algebras ([19, 21, 22]). For the first recall the definition of a lattice ordered group (*l*-group).

**Definition 1.** By an *l*-group  $G$  we consider an algebraic system  $(G, +, \leq)$  such that

1.  $(G, +)$  is commutative group,
2.  $(G, \leq)$  is lattice,
3.  $a \leq b \implies a + c \leq b + c.$

We will use the form of  $MV$ -algebra which is generated by fuzzy sets ([25], [26], [27]). Let  $u \in G, u > 0$ . Then an  $MV$ -algebra  $\mathcal{M}$  is a set  $M = [0, u]$  for which it holds

$$a \oplus b = (a + b) \wedge u,$$

$$a \odot b = (a + b - u) \wedge 0.$$

The element  $u$  is supposed to be a strong unit of  $G$ . Then an  $MV$ -algebra state ([13, 15, 16, 20]) is a mapping  $\bar{m} : \mathcal{M} \rightarrow [0, 1]$  satisfying the following conditions:

1.  $\bar{m}(0) = 0, \bar{m}(u) = 1$ .
2.  $a \odot b = 0 \Rightarrow \bar{m}(a \oplus b) = \bar{m}(a) + \bar{m}(b)$ .
3.  $a_n \nearrow a \Rightarrow \bar{m}(a_n) \nearrow \bar{m}(a)$ .

## 2 Representation of states by probabilities

Let us look at the representation of states from two points of view. The first will be general representation, the second local representation.

### 2.1 General representation of states

Very interesting is the relation between probability measure and state. In [3] there was proved the Butnariu–Klement theorem:

**Theorem 2.** *To any state  $m_{\mathcal{J}} : \mathcal{J} \rightarrow [0, 1]$  there exists a probability measure  $P : \mathcal{S} \rightarrow [0, 1]$  such that*

$$m_{\mathcal{J}}(f) = \int_{\Omega} f dP$$

for any  $f \in \mathcal{J}$ .

This theorem has been generalized also for Atanassov  $IF$ -sets. The generalization of the Butnariu–Klement theorem have the form:

**Theorem 3.** *If  $m : \mathcal{F} \rightarrow [0, 1]$  is a state then there exists a probability measure  $P : \mathcal{S} \rightarrow [0, 1]$  and  $\alpha \in [0, 1]$  such that*

$$m(A) = \int_{\Omega} \mu_A dP + \alpha \left( 1 - \int_{\Omega} (\mu_A + \nu_A) dP \right)$$

for any  $A = (\mu_A, \nu_A) \in \mathcal{F}$ .

This theorem is called Representation theorem for probabilities on  $IF$ -states and it was proved in [4–6, 10, 18]. The Butnariu–Klement theorem can be obtained as a consequence if  $\nu_A = 1 - \mu_A$ .

## 2.2 Local representation of states

In this section we will look at the relations between the observables and random variables. Let start with definitions of these structures [21].

**Definition 4.** An observable is a mapping  $x : \mathcal{B}(R) \rightarrow \mathcal{F}$  satisfying the following conditions

1.  $x(R) = (1_\Omega, 0_\Omega)$ ,
2.  $A \cap B = \emptyset \Rightarrow x(A) \odot x(B) = (0_\Omega, 1_\Omega)$  and  $x(A \cup B) = x(A) \oplus x(B)$ ,
3.  $A_n \nearrow A \Rightarrow x(A_n) \nearrow x(A)$ .

where  $A, A_n, B \in \mathcal{F}$  for  $n = 1, 2, \dots$

**Definition 5.** A random variable is such function  $\xi : \Omega \rightarrow R$  that for any  $A \in \mathcal{B}(R)$  it holds  $\xi^{-1}(A) \in \mathcal{F}$ .

It is easy to prove (see [21]) that:

**Proposition 6.** For any random variable  $\xi : \Omega \rightarrow R$  it holds

1.  $\xi^{-1}(R) = \Omega$ ,
2.  $\xi^{-1}(A \cup B) = \xi^{-1}(A) \cup \xi^{-1}(B)$ ,
3.  $\xi^{-1}(A_n) \nearrow \xi^{-1}(A)$ .

**Proposition 7.** Let  $\Omega = R^N$ . Define the function  $\xi_n : R^N \rightarrow R$  by the formula  $\xi_n((t_i)_{i=1}^\infty) = t_n$ . Then  $\xi$  is the random variable.

Denote by  $\mathcal{C}$  the system of all sets of the form

$$A = \{(t_i)_{i=1}^\infty; t_1 \in A_1, \dots, t_n \in A_n\}$$

where  $n \in N$  and  $A_1, \dots, A_n \in \mathcal{B}(R)$  are Borel sets. To any sequence  $(x_n)_n$  of observables there exist the probability space  $(R^N, \sigma(\mathcal{C}), P)$  such that convergences of  $(x_n)_n$  and  $(\xi_n)_n$  are in correlation. Of course, the space  $(R^N, \sigma(\mathcal{C}), P)$  depends on a concrete sequence  $(x_n)_n$ . For different sequences various spaces can be obtain. We could speak about following types of convergences (for the proof see [19, 21]).

**Proposition 8.** Let  $(x_n)_n$  be a sequence of observables,  $(\xi_n)_n$  be the sequence of corresponding random variables. Then

1.  $(x_n)_n$  converges to  $F : R \rightarrow R$  in distribution if and only if  $(\xi_n)_n$  converges to  $F$ ;
2.  $(x_n)_n$  converges to 0 in state  $m : \mathcal{F} \rightarrow [0, 1]$  if and only if  $(\xi_n)_n$  converges to 0 in measure  $P : \mathcal{S} \rightarrow [0, 1]$ ;
3. if  $(\xi_n)_n$  converges  $P$ -almost everywhere to 0 then  $(x_n)_n$  converges  $m$ -almost everywhere to 0.

### 3 Embedding of fuzzy spaces to $MV$ -algebra

Recall that  $IF$ -set can be geometrically regarded as a function  $A : \Omega \rightarrow \Delta$  where

$$\Delta = \{(x, y); x \geq 0, y \geq 0, x + y \leq 1\}$$

(see *Figure 1*).




Figure 1. Elements of  $IF$ -sets

Let us take  $MV$ -algebra  $\mathcal{M}$  such that  $G = R^2$ . Then the addition of two elements is defined as

$$C = A + B = (\mu_A + \mu_B, \nu_A + \nu_B - 1_\Omega).$$

This addition represents the addition of two vectors with fixed point  $(0, 1)$  (see *Figure 2*). Therefore there is a natural question about the possibility to embedding the  $IF$ -set to  $MV$ -algebra.




Figure 2. Addition in  $MV$ -algebra

For any point  $A = (\mu_A, \nu_A) \in \mathcal{M}$  it holds (see *Figure 3*)

$$(\mu_A, \nu_A) + (0_\Omega, 1_\Omega - \nu_A) = (\mu_A, 0_\Omega).$$




Figure 3. Embedding  $IF$ -set to  $MV$ -algebra

From this reason we can define the function  $\bar{m} : \mathcal{M} \rightarrow [0, 1]$  by the formula

$$\bar{m}(\mu_A, \nu_A) = m(\mu_A, 0_\Omega) - m(0_\Omega, 1_\Omega - \nu_A)$$

This function represent the state on  $\mathcal{M}$  generated by a state defined on  $\mathcal{F}$ .

Recall that there is defined another structure as generalization of fuzzy sets, called interval valued fuzzy set (shortly  $IV$ -set) which was defined by Zadeh ([27]). The  $IV$ -set is a pair of functions  $A = (\mu_A, \nu_A); \mu_A : \Omega \rightarrow [0, 1], \nu_A : \Omega \rightarrow [0, 1]$  for which it holds  $\nu_A \geq \mu_A$ . The ordering on this set is given by

$$A \leq B \Leftrightarrow \mu_A \leq \mu_B, \nu_A \leq \nu_B$$

and the operation  $+$  is defined by

$$A + B = (\mu_A + \mu_B, \nu_A + \nu_B).$$

Also the family  $\mathcal{K}$  of all  $IV$ -sets can be embedded to an  $MV$ -algebra. Here again  $G = R^2$ . Of course the ordering in  $G$  and also the operation  $+$  are given by the same way as in the  $IV$ -sets.




Figure 4.  $IV$ -set

## 4 Review of results

There are many  $IF$ -state applications. We will mention just some of them. In probability theory, there was proved for example central limit theorem [19, 21], law of large numbers [13], martingale convergence theorem [12, 24]. In  $IF$ -dynamical system it was proved Poincare recurrence theorem [20], individual ergodic theorem [19, 21], invariant states [14], entropy of  $IF$ -dynamical system [7–9, 17, 23].

## References

- [1] Atanassov K. T. (1999) *Intuitionistic Fuzzy Sets: Theory and Applications*. Springer Physica-Verlag, Heidelberg.
- [2] Atanassov, K. T. (2012) *On Intuitionistic Fuzzy Set Theory*. Springer, Berlin.
- [3] Butnariu, D. & Klement, E. P. (1993) Triangular norm-based measures. In *Triangular Norm–Based Measures and Games with Fuzzy Coalitions*, 37–68.
- [4] Ciungu, L. & Riečan, B. (2009) General form of probabilities on IF-sets. *Fuzzy Logic and Applications*, 101–107.
- [5] Ciungu, L. C. & Riečan, B. (2010) Representation theorem for probabilities on IFS-events. *Information Sciences*, 180(5), 793–798.
- [6] Čunderlíková, K. & Riečan, B. (2016, October) On Two Formulations of the Representation Theorem for an IF–state. In *International Workshop on Intuitionistic Fuzzy Sets and Generalized Nets*, 65–70.
- [7] Di Nola, A., Dvurečenskij, A., Hyčko, M. & Manara, C. (2005) Entropy on effect algebras with the Riesz decomposition property I: Basic properties, *Kybernetika*, 41(2), 143–160.
- [8] Di Nola, A., Dvurečenskij, A., Hyčko, M. & Manara, C. (2005) Entropy on effect algebras with Riesz decomposition property II: MV-algebras., *Kybernetika*, 41(2), 161–176.
- [9] Ďurica, M. (2007) Entropy on IF–events. entropy. In *Third International Workshop on IFSs, Banská Bystrica, Slovakia, 3 Oct. 2007, Notes on Intuitionistic Fuzzy Sets*, 13(4), 30–40.
- [10] Grzegorzewski, P. & Mrówka, E. (2002) Probability of intuitionistic fuzzy events. In *Soft Methods in Probability, Statistics and Data Analysis*, 105–115.
- [11] Halmos, P. R. (1950) *Measure Theory*, Springer, New York.
- [12] Jurečková, M. (2001) On the conditional expectation on probability MV-algebras with product. *Soft Computing – A Fusion of Foundations, Methodologies and Applications*, 5(5), 381–385.

- [13] Jurečková, M. & Riečan, B. (1997) Weak law of large numbers for weak observables in MV algebras. *Tatra Mountains Math. Publ*, 12, 221–228.
- [14] Michalíková, A. & Riečan, B. (2017) On Invariant Measures on Intuitionistic Fuzzy Sets. In *Advances in Fuzzy Logic and Technology*, 529–534.
- [15] Riečan, B. (1999) On the conditional expectation of observables in MV algebras of fuzzy sets. *Fuzzy Sets and Systems*, 102(3), 445–450.
- [16] Riečan, B. (2000) On the probability theory on MV algebras. *Soft Computing – A Fusion of Foundations, Methodologies and Applications*, 4(1), 49–57.
- [17] Riečan, B. (2005) Kolmogorov – Sinaj entropy on MV–algebras. *International Journal of Theoretical Physics*, 44(7), 1041–1052.
- [18] Riečan, B. (2006) On a problem of Radko Mesiar: General form of IF-probabilities. *Fuzzy Sets and Systems*, 157(11), 1485–1490.
- [19] Riečan B. (2012) Analysis of Fuzzy Logic Models. In: *Intelligent Systems (ed. V. Koleshko) INTECH*, 217 – 244.
- [20] Riečan, B. (2010) Strong Poincaré recurrence theorem in MV-algebras. *Mathematica Slovaca*, 60(5), 655–664.
- [21] Riečan, B. & Mundici, D. (2002) Probability on MV-algebras. *Handbook of Measure Theory*. E. Pap Ed. Elsevier Science, Amsterdam, Chapter 21, 869–910.
- [22] Riečan, B. & Neubrunn, T. (2013) *Integral, Measure, and Ordering*, 411. Springer Science and Business Media.
- [23] Valenčáková, V. (2009) A note on the conditional probability of IF-events. *Mathematica Slovaca*, 59(2), 251–260.
- [24] Vrábelová, M. (2000) A note on the conditional probability on product MV algebras. *Soft Computing*, 4(1), 58–61.
- [25] Zadeh, L. A. (1965) Fuzzy sets, *Information and Control*, 8(3), 338–353.
- [26] Zadeh, L. A. (1968) Probability measures of fuzzy events. *Journal of Mathematical Analysis and Applications*, 23(2), 421–427.
- [27] Zadeh, L. A. (1975) The concept of a linguistic variable and its application to approximate reasoning I. *Information Sciences*, 8(3), 199–249.