

About the Complement in Intuitionistic Fuzzy Sets

H. Bustince, V. Mohedano

Departamento de Automática y Computación.

Public University of Navarre

Abstract: In this paper from the definition of Intuitionistic Fuzzy Sets we analyze the Complementation in these sets. We start by defining the intuitionistic fuzzy complementation in order to then study the particular cases for which this definition coincides with the fuzzy complementation. Afterwards, we analyse the existence of equilibrium points, dual points and we present characterization theorems of intuitionistic fuzzy complements. Lastly, we study a manner of constructing intuitionistic fuzzy sets and analyse the structure of the complementary of intuitionistic fuzzy sets built.

Keywords: Intuitionistic fuzzy set, Intuitionistic fuzzy complementary, fuzzy negation, intuitionistic fuzzy negation.

1. Introduction

Let $X \neq \emptyset$ be a given set. [2] An *intuitionistic fuzzy set* in X is an expression A given by

$$A = \{ \langle x, \mu_A(x), \nu_A(x) \rangle \mid x \in X \}$$

where

$$\mu_A : X \rightarrow [0, 1]$$

$$\nu_A : X \rightarrow [0, 1]$$

with the condition

$$0 \leq \mu_A(x) + \nu_A(x) \leq 1,$$

for all x in X .

The numbers $\mu_A(x)$ and $\nu_A(x)$ denote respectively the degree of membership and the degree of non-membership of the element x in the set A . We will denote $IFSs(X)$ the set of all the

Correspondence to: H. Bustince, Departamento de Automática y Computación, Universidad Pública de Navarra, 31006, Campus Arrosadía, Pamplona, Spain, Phone: 3448169254, Fax: 3448169281, e-mail: bustince@si.upna.es

intuitionistic fuzzy sets in X . Obviously, when

$$\nu_A(x) = 1 - \mu_A(x)$$

for every x in X , the set A is a fuzzy set. We will denote the set of all the fuzzy sets in X with $FSs(X)$.

The following expressions are defined in [2, 3, 6, 7, 8, 9, 10] for every $A, B \in IFSs(X)$

1. $A \leq B$ if and only if $\mu_A(x) \leq \mu_B(x)$ and $\nu_A(x) \geq \nu_B(x)$ for all $x \in X$
2. $A \preceq B$ if and only if $\mu_A(x) \leq \mu_B(x)$ and $\nu_A(x) \leq \nu_B(x)$ for all $x \in X$
3. $A = B$ if and only if $A \leq B$ and $B \leq A$
4. $A_c = \{ \langle x, \nu_A(x), \mu_A(x) \rangle \mid x \in X \}$

We will call

$$\pi_A(x) = 1 - \mu_A(x) - \nu_A(x)$$

intuitionistic index of the element x in the set A .

In 1986, K. Atanassov established different ways of changing an intuitionistic fuzzy set into a fuzzy set and he defined the following operator:

If $A \in IFSs(X)$, then

$$D_\alpha(A) = \{ \langle x, \mu_A(x) + \alpha \cdot \pi_A(x), 1 - \mu_A(x) - \alpha \cdot \pi_A(x) \rangle \mid x \in X \} \quad (1)$$

with $\alpha \in [0, 1]$. Obviously $D_\alpha(A) \in FSs(X)$.

A study of the properties of this operator, (we will call it *Atanassov's operator*), is made in [4, 5].

It is known that if $A' = \{ \langle x, \mu_A(x) \rangle \mid x \in X \}$ is a fuzzy sets on a referential X and $\varphi : [0, 1] \rightarrow [0, 1]$ is a fuzzy complement, the set

$$A = \{ \langle x, \mu_A(x), \varphi(\mu_A(x)) \rangle \mid x \in X \}$$

is not always an intuitionistic fuzzy set. For example, if we take Sugeno's negation [12]

$$\varphi_\lambda(x) = \frac{1-x}{1+\lambda x} \quad \text{with} \quad -1 < \lambda < 0$$

or we take Yager's negation [15, 16]

$$\varphi_\omega(x) = (1-x^\omega)^{\frac{1}{\omega}} \quad \text{with} \quad 1 < \omega$$

the set

$$A = \{ \langle x, \mu_A(x), \varphi(\mu_A(x)) \rangle \mid x \in X \}$$

is not an intuitionistic fuzzy set because

$$\mu_A(x) + \varphi(\mu_A(x)) > 1 \quad \text{for all } x$$

A general study of the fuzzy negations is made in [1, 13, 15].

In this paper we set out to study functions $\varphi : [0, 1] \rightarrow [0, 1]$ such that from a fuzzy set the sum

$$\mu_A(x) + \varphi(\mu_A(x)) \leq 1 \quad \text{for all } x$$

We will conclude the paper studying the conditions in which from a fuzzy set $A' = \{ \langle x, \mu_A(x) \rangle \mid x \in X \}$ we build an intuitionistic fuzzy set $A = \{ \langle x, \mu_A(x), \nu_A(x) \rangle \mid x \in X \}$.

2. Intuitionistic Fuzzy Complement

Definition 1. A function $\varphi : [0, 1] \rightarrow [0, 1]$ will be called *intuitionistic fuzzy complement* iff

$$\varphi(x) \leq 1 - x, \quad \text{for all } x \in [0, 1].$$

An intuitionistic fuzzy complement will be called *continuous*, *decreasing*, *increasing* iff φ is continuous, decreasing, increasing respectively.

Note that according to the definition above, $\varphi(0) \leq 1$ and $\varphi(1) = 0$.

Hereinafter we will denote the intuitionistic fuzzy complement IFC.

We have to keep in mind that there are fuzzy complements that are not IFC, for example Sugeno's

$$\varphi_\lambda(x) = \frac{1 - x}{1 + \lambda \cdot x}$$

for $-1 < \lambda < 0$, Yager's complement [15, 16]

$$\varphi_\omega(x) = (1 - x^\omega)^{\frac{1}{\omega}}$$

for $\omega > 1$.

Obviously, if we impose the conditions:

i) $\varphi(0) = 1$

ii) φ is non-increasing

on an intuitionistic fuzzy complement, we have a fuzzy negation.

Proposition 1. *The only continuous intuitionistic fuzzy complement constant for all $x \in [0, 1]$ is the intuitionistic fuzzy complement $\varphi(x) = 0$ for all $x \in [0, 1]$.*

Proof. Evident. $\square$

Definition 2. Let us say that $x \in [0, 1]$ is an *equilibrium point* for the intuitionistic fuzzy complement φ iff $\varphi(x) = x$.

Theorem 1. *Every continuous intuitionistic fuzzy complement has at most one equilibrium point.*

Proof. Let φ be a continuous intuitionistic fuzzy complement. An equilibrium point is a solution of the equation $\varphi(x) - x = 0$, where $x \in [0, 1]$. Let us suppose that $\varphi(x) - x = b$, where b is a real number, and we take $F(x) = \varphi(x) - x = b$. As φ is an intuitionistic fuzzy complement we have $F(1) = \varphi(1) - 1 = -1$ and $F(0) = \varphi(0) \leq 0$. If $F(0) > 0$ then, by Rolle's theorem there is at least a $c \in [0, 1]$ such that $F(c) = 0$, therefore $\varphi(c) = c$. If $\varphi(0) = 0$, obviously we have that the 0 is a point of equilibrium. $\square$

Theorem 2. *Let $\varphi : [0, 1] \rightarrow [0, 1]$ an intuitionistic fuzzy complementation.*

- i) *If $x \in [0, 1]$ is an equilibrium point, then $x \leq 1/2$.*
- ii) *If φ is continuous and decreasing in $[0, 1]$, then φ has a unique equilibrium point.*

Proof. i) $\varphi(x) = x \leq 1 - x$, then $x \leq 1/2$.

ii) Let us suppose that it is not true that there is a unique equilibrium point, that is, there are $x_1, x_2 \in [0, 1]$ such that $x_1 < x_2$ and $\varphi(x_1) = x_1$ and $\varphi(x_2) = x_2$. Then, as φ is decreasing, we have that if $x_1 < x_2$, then $\varphi(x_1) \geq \varphi(x_2)$, therefore $x_1 \geq x_2$ in contradiction with the hypothesis that $x_1 < x_2$, so $x_1 = x_2$.

Definition 3. Let $\varphi : [0, 1] \rightarrow [0, 1]$ be an intuitionistic fuzzy complementation, and let $x, y \in [0, 1]$. We will say that x is the *dual point* of y with respect to φ or that y is the *dual point* of x with respect to φ if $\varphi(x) + \varphi(y) = x + y$ holds.

Theorem 3. *Let $\varphi : [0, 1] \rightarrow [0, 1]$ an intuitionistic fuzzy complementation.*

- i) *If x is the dual point of y with respect to φ , then $x + y \leq 1$.*
- ii) *The number 1 has dual point iff $\varphi(0) = 1$*
- iii) *Every equilibrium point is the dual point its self.*
- iv) *Each pair of equilibrium point of φ are dual points between each other.*

Proof. i) $x + y = \varphi(y) + \varphi(x) \leq 1 - y + 1 - x = 2 - (x + y)$, then $x + y \leq 1$.

ii) $\Rightarrow$ $\varphi(1) + \varphi(y) = 1 + y$, as $\varphi(1) = 0$ then $\varphi(y) = 1 + y \leq 1$, therefore $y = 0$, then $\varphi(0) = 1$.

$\Leftarrow$ If $\varphi(0) = 1$, then $\varphi(0) + \varphi(1) = 1 + 0$.

iii) Evident.

iv) Let $\varphi(x) = x$ and $\varphi(y) = y$, then $\varphi(x) + \varphi(y) = x + y$. $\square$

Definition 3. An intuitionistic fuzzy complement, $\varphi : [0, 1] \rightarrow [0, 1]$, will be called *involution* in $[0, p]$, with $p \in [0, 1]$ iff $\varphi(\varphi(x)) = x$ for all $x \in [0, p]$.

Theorem 4. Let $\varphi : [0, 1] \rightarrow [0, 1]$ be an intuitionistic fuzzy complementation. Then, $\varphi(x)$ is the dual point of x for all $x \in [0, p]$ with $p \in [0, 1]$ iff φ is involutive in $[0, p]$.

Proof. $\Rightarrow$) As $\varphi(x)$ is the dual point of x for all $x \in [0, p]$ we have that $\varphi(\varphi(x)) + \varphi(x) = x + \varphi(x)$ for all $x \in [0, p]$.

$\Leftarrow$) As φ is involutive in $[0, p]$ we have that for all $x \in [0, p]$, $\varphi(\varphi(x)) = x$, then $\varphi(\varphi(x)) + \varphi(x) = x + \varphi(x)$. $\square$

Theorem 5. Let $\varphi : [0, 1] \rightarrow [0, 1]$ be an intuitionistic fuzzy complementation continuous, strictly decreasing and involutive in $[0, p]$ with $p \in [0, 1]$ and $\varphi(x) = 0$ for all $x \in [p, 1]$. Then $\varphi(0) = p$.

Proof. As $p = \varphi(\varphi(p)) = \varphi(0)$. $\square$

Obviously, in the same conditions as the previous theorem $\varphi^{-1} = \varphi$ in $[0, p]$.

3. Characterization Theorems of Intuitionistic Fuzzy Complements

In this section we will denote the standard negation N , $N : [0, 1] \rightarrow [0, 1]$ given by $N(x) = 1 - x$ for all $x \in [0, 1]$.

Theorem 6. Let $\varphi : [0, 1] \rightarrow [0, 1]$. Then,

φ is a continuous intuitionistic fuzzy complement if and only if there exists a continuous function $f : [0, 1] \rightarrow [0, 1]$ such that

- i) $f(x) \leq x$ for all $x \in [0, 1]$
- ii) $\varphi(x) = (f \circ N)(x)$ for all $x \in [0, 1]$.

Proof. $\Rightarrow$) φ is a continuous intuitionistic fuzzy complement. We will consider the function $f(x) = \varphi(1 - x)$ for all $x \in [0, 1]$. Evidently, φ is continuous, besides $f(x) = \varphi(1 - x) \leq x$ for all $x \in [0, 1]$. We can say that f satisfies iii), $(f \circ N)(x) = f(N(x)) = f(1 - x) = \varphi(x)$ for all $x \in [0, 1]$.

$\Leftarrow$) $\varphi(x) = (f \circ N)(x) = f(N(x)) = f(1 - x) \leq 1 - x$ for all $x \in [0, 1]$. Evidently φ is a continuous function. $\square$

An example of this theorem is the following:

$$f(x) = \begin{cases} \frac{1}{2}x & \text{if } x \leq \frac{1}{4} \\ \frac{1}{6} - \frac{1}{6}x & \text{if } x \geq \frac{1}{4} \end{cases}$$

the corresponding IFN is

$$\varphi(x) = \begin{cases} \frac{1}{6}x & \text{if } x \leq \frac{3}{4} \\ \frac{1}{2}(1 - x) & \text{if } x \geq \frac{3}{4} \end{cases}$$

Theorem 7. Let $\varphi : [0, 1] \rightarrow [0, 1]$. Then,

φ is a continuous, monotone decreasing intuitionistic fuzzy complement if and only if there exists a continuous function $f : [0, 1] \rightarrow [0, 1]$ such that

- i) monotone increasing
- ii) $f(x) \leq x$ for all $x \in [0, 1]$
- iii) $\varphi(x) = (f \circ N)(x)$ for all $x \in [0, 1]$.

Proof. $\Rightarrow$), Similar to above, we will now say that f is increasing, let $x_1, x_2 \in [0, 1]$ such that $x_1 < x_2$, as φ is decreasing we have $\varphi(x_1) \geq \varphi(x_2)$, therefore $f(1 - x_1) \geq f(1 - x_2)$ and besides $1 - x_1 \geq 1 - x_2$ we have that f is increasing.

$\Leftarrow$) Similar to what was done previous theorem, we will now see that φ is decreasing, let $x_1, x_2 \in [0, 1]$ such that $x_1 < x_2$, therefore $1 - x_1 > 1 - x_2$ as f is increasing we have that $f(1 - x_1) \geq f(1 - x_2)$ therefore $\varphi(x_1) \geq \varphi(x_2)$. $\square$

An example of this theorem is the following:

$$f(x) = \begin{cases} 0 & \text{if } x \leq \frac{1}{4} \\ (\frac{2}{3}x - \frac{1}{6})^2 & \text{if } x \geq \frac{1}{4} \end{cases}$$

the corresponding IFC is

$$\varphi(x) = \begin{cases} 0 & \text{if } x \geq \frac{3}{4} \\ (\frac{1}{2} - \frac{2}{3}x)^2 & \text{if } x \leq \frac{3}{4} \end{cases}$$

6. Construction of an IFSs from a FSs.

In this section we present a method of construction of intuitionistic fuzzy sets from fuzzy sets. This method is basically constructing an IFSs from the membership functions of the fuzzy set considered and the intuitionistic fuzzy complement analysed in the sections above. Afterwards we will see the general structure of the complementaries of the fuzzy sets built.

Theorem 8. Let $A' = \{ \langle x, \mu_A(x) \rangle \mid x \in X \}$ be an fuzzy set on the referential set $X \neq \emptyset$, and let φ an intuitionistic fuzzy complementation. The set

$$\{ \langle x, \mu_A(x), \varphi(\mu_A(x)) \rangle \mid x \in X \}$$

is an intuitionistic fuzzy set on X .

Proof. We only need to recall definition 1. $\square$

We will denote the intuitionistic fuzzy sets built in the way indicated in the theorem above $\varphi(A)$.

It is clear that from functions f that satisfy the conditions of the theorems exposed in the section above we can build intuitionistic fuzzy complements that when applied to the membership functions of fuzzy sets we get intuitionistic fuzzy sets satisfying the conditions that we have imposed on the functions f .

It is interesting to note that intuitionistic index of each element of this set is the following: $\pi_{\varphi(A)}(x) = 1 - \mu_A(x) - \varphi(\mu_A(x))$, obviously $0 \leq \pi_{\varphi(A)}(x) \leq 1$ for all $x \in X$. Besides, taking into account the definition of D_α , we have that

$$D_0(\varphi(A)) = A' = \{ \langle x, \mu_A(x) \rangle \mid x \in X \}$$

$$D_1(\varphi(A)) = \{ \langle x, 1 - \varphi(\mu_A(x)) \rangle \mid x \in X \}$$

It is interesting to point out particular case in which the functions f of the theorems of the section above coincide with the membership function μ .

Corollary 1.

Let $A' \in FSS(X)$ such that $\mu_A : [0, 1] \rightarrow [0, 1]$ continuous.

If $\mu_A(x) \leq x$ for all $x \in [0, 1]$, then $\{ \langle x, \mu_A(x), \nu_A(x) = \mu_A(1 - x) \rangle \mid x \in [0, 1] \}$ is an intuitionistic fuzzy set.

7. References

- [1] C. Alsina, E. Trillas, L. Valverde, On some logical connectives for Fuzzy Sets Theory, J. Math. An. and Appl., 93 (1983) 15-26.
- [2] K. Atanassov, Intuitionistic fuzzy sets, Fuzzy sets and Systems, 20 (1986) 87-96.
- [3] K. Atanassov, Review and New Results on Intuitionistic Fuzzy Sets, IM-MFAIS, 1 (1988).
- [4] K. Atanassov, Two operators on intuitionistic fuzzy sets, Comptes rendus de l'Académie bulgare des Sciences, Tome 41, 5 (1988).
- [5] P. Burillo, H. Bustince, Construction theorems for intuitionistic fuzzy sets, will be published in Fuzzy sets and Systems, 84 (1996)
- [6] P. Burillo, H. Bustince, Entropy on intuitionistic fuzzy sets and on interval-valued fuzzy sets, Fuzzy sets and Systems, 78 (1996) 305-316.
- [7] P. Burillo, H. Bustince, Intuitionistic fuzzy relations (Part I), Mathware and Soft Computing, 2 (1995) 5-38.
- [8] P. Burillo and H. Bustince, Estructuras algebraicas en conjuntos IFS, II Congreso Nacional de Lógica y Tecnología Fuzzy, Boadilla del Monte, Madrid, Spain (1992) 135-147.
- [9] H. Bustince, Conjuntos Intuicionistas e Intervalo valorados Difusos: Propiedades y Construcción. Relaciones Intuicionistas Fuzzy. Thesis, Universidad Pública de Navarra (1994).

- [10] H. Bustince, P. Burillo, Correlation of interval-valued intuitionistic fuzzy sets, *Fuzzy sets and Systems*, 74 (1995) 237-244.
- [11] H. Bustince, P. Burillo, Structures on intuitionistic fuzzy relations, *Fuzzy sets and Systems*, 78 (1996) 293-303.
- [12] M. Sugeno, Theory of fuzzy integrals and its applications. Thesis. Tokyo Institute of Technology (1974).
- [13] E. Trillas, Sobre funciones de negacion en la teoria de conjuntos difusos, *Stochastica*, III-1 (1979) 47-59.
- [14] E. Trillas, C. Alsina, J.M. Terricabras, Introducci3n a la l3gica borrosa, *Ariel Matemática*, (1995).
- [15] R. Yager, On a general class of fuzzy connectives, *Fuzzy Sets and Systems*, 4 (1980) 338-353.
- [16] R. Yager, On the measure of fuzziness and negation. Part I: Membership in the unit interval, *Intern. J. of General Systems*, 5 (1979) 221-229.