

ICIFSS, 8–10 January 2018, Erode, Tamil Nadu, India

Notes on Intuitionistic Fuzzy Sets

Print ISSN 1310–4926, Online ISSN 2367–8283

Vol. 24, 2018, No. 1, 37–47

DOI: 10.7546/nifs.2018.24.1.37-47

# The Cauchy problem for intuitionistic fuzzy differential equations

**Bouchra Ben Amma, Said Melliani and Lalla Saadia Chadli**

Laboratory of Applied Mathematics and Scientific Computing (LMACS)

Sultan Moulay Slimane University

PO Box 523, 23000 Beni Mellal, Morocco

e-mails: bouchrabenamma@gmail.com, s.melliani@yahoo.fr,

sa.chadli@yahoo.fr

**Received:** 20 October 2017

**Revised:** 4 December 2017

**Accepted:** 6 December 2017

**Abstract:** In this paper we discuss the existence and uniqueness theorem of a solution of the cauchy problem of intuitionistic fuzzy differential equation.

**Keywords:** Intuitionistic fuzzy differential equations, Approximate solutions, Existence, Uniqueness.

**2010 Mathematical Subject Classification:** 03E72.

## 1 Introduction

One of the generalizations of fuzzy sets theory [15] can be considered the proposed intuitionistic fuzzy sets (IFS). Later on Atanassov generalized the concept of fuzzy set and introduced the idea of intuitionistic fuzzy set [1–3]. They are very necessary and powerful tool in modeling imprecision, valuable applications of IFSs have been flourished in many different field [4, 6, 8, 9].

For intuitionistic fuzzy concepts, recently the authors [5, 11–13] established, the theory of metric space of intuitionistic fuzzy sets, intuitionistic fuzzy differential equations, intuitionistic fuzzy fractional equation and the Cauchy problem for complex intuitionistic fuzzy differential equations. They proved the existence and uniqueness of the intuitionistic fuzzy solution for these intuitionistic fuzzy differential equations using different concepts.

This paper is to investigate the existence and uniqueness theorem of intuitionistic fuzzy solutions for the following intuitionistic fuzzy differential equations:

$$(\langle u, v \rangle)'(t) = f(t, \langle u, v \rangle(t)), \quad \langle u, v \rangle(t_0) = \langle u_0, v_0 \rangle \quad (1.1)$$

when  $\langle u_0, v_0 \rangle$  is an intuitionistic fuzzy quantity and  $f$  satisfies the generalized Lipschitz condition.

The paper is organized as follows. In Section 2, we collect the fundamental notions and facts which will be used in the rest of the article and we list several comparison propositions on classical ordinary differential equations in [7]. In Section 3 we show the relation between a solution and its approximate solution to the Cauchy problem of the intuitionistic fuzzy differential equation, and furthermore, in Section 4, we prove the existence and uniqueness theorem for a solution to the Cauchy problem of the intuitionistic fuzzy differential equation.

## 2 Preliminaries

Throughout this paper,  $(\mathbb{R}^n, B(\mathbb{R}^n), \mu)$  denotes a complete finite measure space.

Let us  $P_k(\mathbb{R}^n)$  the set of all non empty compact convex subsets of  $\mathbb{R}^n$ . we denote by

$$\mathbb{F}_n = \text{IF}(\mathbb{R}^n) = \{ \langle u, v \rangle : \mathbb{R}^n \rightarrow [0, 1]^2, |\forall x \in \mathbb{R}^n \ 0 \leq u(x) + v(x) \leq 1 \}$$

An element  $\langle u, v \rangle$  of  $\mathbb{F}_n$  is said an intuitionistic fuzzy number if it satisfies the following conditions

- (i)  $\langle u, v \rangle$  is normal i.e there exists  $x_0, x_1 \in \mathbb{R}^n$  such that  $u(x_0) = 1$  and  $v(x_1) = 1$ .
- (ii)  $u$  is fuzzy convex and  $v$  is fuzzy concave.
- (iii)  $u$  is upper semi-continuous and  $v$  is lower semi-continuous
- (iv)  $\text{supp } \langle u, v \rangle = \text{cl}\{x \in \mathbb{R}^n : |v(x) < 1\}$  is bounded.

so we denote the collection of all intuitionistic fuzzy numbers by  $\mathbb{F}_n$

On the space  $\mathbb{F}_n$  we will consider the following metric,

$$\begin{aligned} d_\infty^n \left( \langle u, v \rangle, \langle z, w \rangle \right) &= \frac{1}{4} \sup_{0 < \alpha \leq 1} \left\| \left[ \langle u, v \rangle \right]_r^+(\alpha) - \left[ \langle z, w \rangle \right]_r^+(\alpha) \right\| \\ &+ \frac{1}{4} \sup_{0 < \alpha \leq 1} \left\| \left[ \langle u, v \rangle \right]_l^+(\alpha) - \left[ \langle z, w \rangle \right]_l^+(\alpha) \right\| + \frac{1}{4} \sup_{0 < \alpha \leq 1} \left\| \left[ \langle u, v \rangle \right]_r^-(\alpha) - \left[ \langle z, w \rangle \right]_r^-(\alpha) \right\| \\ &+ \frac{1}{4} \sup_{0 < \alpha \leq 1} \left\| \left[ \langle u, v \rangle \right]_l^-(\alpha) - \left[ \langle z, w \rangle \right]_l^-(\alpha) \right\| \end{aligned}$$

where  $\| \cdot \|$  denotes the usual Euclidean norm in  $\mathbb{R}^n$ .

**Theorem 2.1** ([12]).  $d_\infty^n$  define a metric on  $\text{IF}_n$ .

**Theorem 2.2** ([12]). The metric space  $(\text{IF}_n, d_\infty^n)$  is complete.

The norm  $\| \cdot \|$  of an intuitionistic fuzzy number  $\langle u, v \rangle \in IF_n$  is defined by

$$\| \langle u, v \rangle \| = d_{\infty}^n(0_{(1,0)}, \langle u, v \rangle) = \| [\langle u, v \rangle]_0 \| = \frac{1}{2} \sup_{a \in [\langle u, v \rangle]_0} |a| + \frac{1}{2} \inf_{b \in [\langle u, v \rangle]_0} |b|$$

**Definition 2.1.** An intuitionistic fuzzy set  $\langle u, v \rangle$  is called convex intuitionistic fuzzy set if and only if  $u$  is convex fuzzy set and  $v$  is concave fuzzy set.

The question that arises, is what  $IF_n$  with addition and multiplication by a scalar is a vector

**Theorem 2.3** ([13]). There exists a normed space  $X$  and a function  $j : IF_n \rightarrow X$  with properties:

1.  $j$  is an isometry i.e.  $\| j(\langle u, v \rangle) - j(\langle u', v' \rangle) \| = d_{\infty}^n(\langle u, v \rangle, \langle u', v' \rangle)$
2.  $j(\langle u, v \rangle \oplus \langle u', v' \rangle) = j(\langle u, v \rangle) + j(\langle u', v' \rangle)$
3.  $j(\lambda \langle u, v \rangle) = \lambda j(\langle u, v \rangle) \quad \lambda \geq 0$

**Remark 2.1.** if  $\langle u, v \rangle(t) : T \rightarrow IF_n$  is differentiable at  $t_0 \in T$ , then

$$(j\langle u, v \rangle)(t) = j(\langle u, v \rangle(t)) \quad T \rightarrow X$$

is Frechet differentiable at  $t_0$  and  $(j\langle u, v \rangle)'(t_0) = j(\langle u, v \rangle'(t_0))$ , where  $j$  is the embedding in Theorem 2.3.

In the following we list several comparison propositions on classical ordinary differential equations following [7]

**Proposition 2.1.** Let  $G \subset \mathbb{R}^2$  be an open set and  $g \in C[G, \mathbb{R}]$ ,  $(t_0, x_0) \in G$ . Suppose  $r(t)$  is the maximum solution to the initial value problem

$$x' = g(t, x), \quad x(t_0) = x_0 \tag{2.1}$$

and its largest interval of existence of right solution is  $[t_0, t_0 + a)$ . If  $[t_0, t_1] \subset [t_0, t_0 + a)$ , then there exists an  $\varepsilon_0 > 0$  such that the maximum solution  $r(t, \varepsilon)$  to the initial value problem

$$x' = g(t, x) + \varepsilon, \quad x(t_0) = x_0 + \varepsilon$$

exists on  $[t_0, t_1]$  whenever  $0 < \varepsilon < \varepsilon_0$ , and  $r(t, \varepsilon)$  uniformly converges to  $r(t)$  on  $[t_0, t_1]$  as  $\varepsilon \rightarrow 0^+$

**Proposition 2.2.** Let  $G \subset \mathbb{R}^2$  be an open set,  $g \in C[G, \mathbb{R}]$ ,  $(t, x_0) \in G$ . Suppose that the maximum solution to the initial value problem (2.1) is  $r(t)$  and its largest interval of existence of right solution is  $[t_0, t_0 + a)$ . If  $m(t) \in C[[t_0, t_0 + a), \mathbb{R}]$ , satisfies  $(t, m(t)) \in G$  for all  $t \in [t_0, t_0 + a)$ ,  $m(t_0) \leq x_0$ , and

$$Dm(t) \leq g(t, m(t)), \quad \forall t \in [t_0, t_0 + a) \setminus \Gamma$$

where  $D$  is one of the four Dini derivatives (see [7]),  $G$  at most is a countable set on  $t$ . Then we must have

$$m(t) \leq r(t), \quad \forall t \in [t_0, t_0 + a)$$

### 3 The relation between a solution and its approximate solution to intuitionistic fuzzy differential equations

Assume that  $f : T \times W \longrightarrow IF_n$  is continuous (it is denoted by  $f \in C[T \times W, IF_n]$ ). Consider the initial value problem

$$\langle u, v \rangle'(t) = f(t, \langle u, v \rangle(t)), \quad \langle u, v \rangle(t_0) = \langle u, v \rangle_0 \quad (3.1)$$

where  $W \subset IF_n$ ,  $\langle u, v \rangle(t_0) \in W$ .

In the following we give the relation between a solution and its approximate solutions. We denote  $R_0 = [t_0, t_0 + p] \times B(\langle u, v \rangle_0, q)$  where  $p > 0$ ,  $q > 0$ ,  $\langle u, v \rangle_0 \in IF_n$ ,  $B(\langle u, v \rangle_0, q) = \{\langle u, v \rangle \in IF_n \mid d_\infty^n(\langle u, v \rangle, \langle u, v \rangle_0) \leq q\}$

**Theorem 3.1.** Let  $f \in C[R_0, IF_n]$ ,  $r \in (0, p)$ ,  $\langle u, v \rangle_n \in C^1[[t_0, t_0 + r], B(\langle u, v \rangle_0, q)]$  such that

$$\begin{aligned} j\langle u, v \rangle'_n(t) &= jf(t, \langle u, v \rangle_n(t)) + B_n(t), \quad \langle u, v \rangle(t_0) = \langle u, v \rangle_0, \\ \|B_n(t)\| &\leq \varepsilon_n \quad \forall t \in [t_0, t_0 + r] \quad (n = 0, 1, 2, \dots), \end{aligned} \quad (3.2)$$

where  $\varepsilon_n > 0$ ,  $\varepsilon_n \longrightarrow 0$ ,  $B_n(t) \in C[[t_0, t_0 + r], X]$  and  $j$  is the isometric embedding from  $(IF_n, d_\infty^n)$  onto its range in the Banach space  $X$ . For each  $t \in [t_0, t_0 + r]$  there exist an  $\delta(t) > 0$  such that  $H$ -differences  $\langle u, v \rangle_n(t+h) \ominus \langle u, v \rangle_n(t)$  and  $\langle u, v \rangle_n(t) \ominus \langle u, v \rangle_n(t-h)$  exist for all  $0 \leq h < \delta(t)$  and  $n = 1, 2, \dots$

if we have

$$d_\infty^n(\langle u, v \rangle_n(t), \langle u, v \rangle(t)) \longrightarrow 0 \quad \text{u.c.} \quad \forall t \in [t_0, t_0 + r] \quad (n \longrightarrow \infty) \quad (3.3)$$

(u.c. denotes the uniform convergence), then  $\langle u, v \rangle \in C^1[[t_0, t_0 + r], B(\langle u, v \rangle_0, q)]$

$$\langle u, v \rangle'(t) = f(t, \langle u, v \rangle(t)), \quad \langle u, v \rangle(t_0) = \langle u_0, v_0 \rangle, \quad t \in [t_0, t_0 + r]. \quad (3.4)$$

*Proof 1.* From (3.3) we know that  $\langle u, v \rangle \in C[[t_0, t_0 + r], B(x_0, q)]$ . For fixed  $t_1 \in [t_0, t_0 + r]$  and any  $t \in [t_0, t_0 + r]$ ,  $t > t_1$ , denote

$$F(t, n) = \frac{j\langle u, v \rangle_n(t) - j\langle u, v \rangle_n(t_1)}{t - t_1} - jf(t_1, \langle u, v \rangle_n(t_1)) - B_n(t_1)$$

It is well known that

$$\lim_{n \rightarrow \infty} F(t, n) = \frac{j\langle u, v \rangle(t) - j\langle u, v \rangle(t_1)}{t - t_1} - jf(t_1, \langle u, v \rangle(t_1)) \quad (3.5)$$

From  $f \in C^1[R_0, IF_n]$  is known that for any  $\varepsilon > 0$ , there exists  $\delta_1 > 0$  such that

$$d_\infty^n\left(f(t, \langle z, w \rangle(t)), f(t_1, \langle u, v \rangle(t_1))\right) < \frac{\varepsilon}{4} \quad (3.6)$$

whenever  $t_1 < t < t_1 + \delta_1$  and  $d_\infty^n(\langle z, w \rangle(t), \langle u, v \rangle(t_1)) < \delta_1$  with  $\langle z, w \rangle \in B(\langle u, v \rangle_0, q)$ .

Take natural number  $N > 0$  such hat

$$\varepsilon_n < \frac{\varepsilon}{4}, \quad d_\infty^n(\langle u, v \rangle_n(t), \langle u, v \rangle(t)) < \frac{\delta_1}{2} \quad \text{for any } n > N, t \in [t_0, t_0 + r]. \quad (3.7)$$

Take  $\delta > 0$  such that  $\delta < \delta_1$  and

$$d_\infty^n \left( \langle u, v \rangle(t), \langle u, v \rangle(t_1) \right) < \frac{\delta_1}{2} \quad \text{whenever } t_1 < t < t_1 + \delta. \quad (3.8)$$

By the definition of  $F(t, n)$  and (3.2), we have

$$j\langle u, v \rangle_n(t) - j\langle u, v \rangle_n(t_1) - (t - t_1)j\langle u, v \rangle'_n(t_1) = (t - t_1)F(t, n) \quad (3.9)$$

We choose  $\varphi \in X^*$  such that  $\|\varphi\| = 1$  and

$$\begin{aligned} & \varphi(j\langle u, v \rangle_n(t) - j\langle u, v \rangle_n(t_1) - (t - t_1)j\langle u, v \rangle'_n(t_1)) \\ & = \|\varphi(j\langle u, v \rangle_n(t) - j\langle u, v \rangle_n(t_1) - (t - t_1)j\langle u, v \rangle'_n(t_1))\| \end{aligned}$$

Let  $\psi(t) = \varphi(j\langle u, v \rangle_n(t)) - (t - t_1)\varphi(j\langle u, v \rangle'_n(t_1))$ , consequently,

$$\psi'(t) = \varphi(j\langle u, v \rangle'_n(t)) - \varphi(j\langle u, v \rangle'_n(t_1))$$

hence

$$\begin{aligned} & \|j\langle u, v \rangle_n(t) - j\langle u, v \rangle_n(t_1) - (t - t_1)j\langle u, v \rangle'_n(t_1)\| = \psi(t) - \psi(t_1) = \psi'(\bar{t})(t - t_1) \\ & = \varphi(j\langle u, v \rangle'_n(\bar{t}) - j\langle u, v \rangle'_n(t_1))(t - t_1) \leq \|\varphi\| \cdot \|j\langle u, v \rangle'_n(\bar{t}) - j\langle u, v \rangle'_n(t_1)\| \cdot (t - t_1) \\ & = \|j\langle u, v \rangle'_n(\bar{t}) - j\langle u, v \rangle'_n(t_1)\| \cdot (t - t_1) \end{aligned}$$

where  $t_1 \leq \bar{t} \leq t$ . In view of (3.9), we have

$$\|F(t, n)\| \leq \|j\langle u, v \rangle'_n(\bar{t}) - j\langle u, v \rangle'_n(t_1)\|, \quad t_1 \leq \bar{t} \leq t \quad (3.10)$$

From (3.7) and (3.8) we know that

$$d_\infty^n \left( \langle u, v \rangle(\bar{t}), \langle u, v \rangle(t_1) \right) < \frac{\delta_1}{2}$$

$$\begin{aligned} d_\infty^n \left( \langle u, v \rangle_n(\bar{t}), \langle u, v \rangle(t_1) \right) & \leq d_\infty^n \left( \langle u, v \rangle_n(\bar{t}), \langle u, v \rangle(\bar{t}) \right) + d_\infty^n \left( \langle u, v \rangle(\bar{t}), \langle u, v \rangle(t_1) \right) \\ & < \frac{\delta_1}{2} + \frac{\delta_1}{2} = \delta_1 \end{aligned}$$

Hence by (3.6) and (3.10) we have

$$\begin{aligned} \|F(t, n)\| & \leq \|j\langle u, v \rangle'_n(\bar{t}) - j\langle u, v \rangle'_n(t_1)\| \\ & = \|jf(\bar{t}, \langle u, v \rangle_n(\bar{t})) + B_n(\bar{t}) - jf(t_1, \langle u, v \rangle_n(t_1)) - B_n(t_1)\| \\ & \leq \|jf(\bar{t}, \langle u, v \rangle_n(\bar{t})) - jf(t_1, \langle u, v \rangle(t_1))\| \\ & + \|jf(t_1, \langle u, v \rangle(t_1)) - jf(t_1, \langle u, v \rangle_n(t_1))\| + 2\varepsilon_n \\ & = d_\infty^n \left( f(\bar{t}, \langle u, v \rangle_n(\bar{t})), f(t_1, \langle u, v \rangle(t_1)) \right) \\ & + d_\infty^n \left( f(t_1, \langle u, v \rangle(t_1)), f(t_1, \langle u, v \rangle_n(t_1)) \right) + 2\varepsilon_n \\ & < \frac{\varepsilon}{4} + \frac{\varepsilon}{4} + 2\varepsilon_n < \varepsilon \end{aligned}$$

whenever  $n > N$  and  $t_1 < t < t_1 + \delta$ .

Now let  $n \rightarrow \infty$ , and applying Eq. (3.5), we have

$$\left\| \frac{j\langle u, v \rangle(t) - j\langle u, v \rangle(t_1)}{t - t_1} - jf(t_1, \langle u, v \rangle(t_1)) \right\| \leq \varepsilon, \quad t_1 < t < t_1 + \delta. \quad (3.11)$$

On the other hand, from the assumption of Theorem 3.1, there exists an  $\delta(t_1) \in (0, \delta)$  such that the H-differences

$$\langle u, v \rangle_n(t) \ominus \langle u, v \rangle_n(t_1)$$

exist for all  $t \in [t_1, t_1 + \delta(t_1)]$  and  $n = 1, 2, \dots$

Let  $\langle z, w \rangle_n(t) = \langle u, v \rangle_n(t) \ominus \langle u, v \rangle_n(t_1)$ . We verify that the intuitionistic fuzzy number-valued sequence  $\{\langle z, w \rangle_n(t)\}$  uniformly converges on  $[t_1, t_1 + \delta(t_1)]$

In fact, from the assumption  $d_\infty^n(\langle u, v \rangle_n(t), \langle u, v \rangle(t)) \rightarrow 0$ , u.c  $\forall t \in [t_0, t_0 + r]$ , we know

$$\begin{aligned} d_\infty^n(\langle z, w \rangle_n(t), \langle z, w \rangle_m(t)) &= d_\infty^n(\langle z, w \rangle_n(t) + \langle u, v \rangle_n(t_1), \langle z, w \rangle_m(t) + \langle u, v \rangle_n(t_1)) \\ &\leq d_\infty^n(\langle u, v \rangle_n(t), \langle u, v \rangle_m(t)) \\ &\quad + d_\infty^n(\langle u, v \rangle_m(t), \langle z, w \rangle_m(t) + \langle u, v \rangle_n(t_1)) \\ &= d_\infty^n(\langle u, v \rangle_n(t), \langle u, v \rangle_m(t)) \\ &\quad + d_\infty^n(\langle z, w \rangle_m(t) + \langle u, v \rangle_m(t_1), \langle z, w \rangle_m(t) + \langle u, v \rangle_n(t_1)) \\ &= d_\infty^n(\langle u, v \rangle_n(t), \langle u, v \rangle_m(t)) + d_\infty^n(\langle u, v \rangle_m(t_1), \langle u, v \rangle_n(t_1)) \\ &\rightarrow 0 \text{ u.c.} \quad \forall t \in [t_1, t_1 + \delta(t_1)] (n, m \rightarrow \infty) \end{aligned}$$

Since  $(IF_n, d_\infty^n)$  is complete, there exists an intuitionistic fuzzy number-valued mapping such that  $\{\langle z, w \rangle_n(t)\}$  uniformly converges to  $\langle z, w \rangle(t)$  on  $[t_1, t_1 + \delta(t_1)]$  as  $n \rightarrow \infty$

In addition, we have

$$\begin{aligned} d_\infty^n(\langle u, v \rangle(t_1) + \langle z, w \rangle(t), \langle u, v \rangle(t)) &\leq d_\infty^n(\langle u, v \rangle(t_1) + \langle z, w \rangle(t), \langle u, v \rangle_n(t_1) + \langle z, w \rangle_n(t)) \\ &\quad + d_\infty^n(\langle u, v \rangle_n(t_1) + \langle z, w \rangle_n(t), \langle u, v \rangle(t)) \\ &\leq d_\infty^n(\langle u, v \rangle(t_1) + \langle z, w \rangle(t), \langle u, v \rangle(t_1) + \langle z, w \rangle_n(t)) \\ &\quad + d_\infty^n(\langle u, v \rangle(t_1) + \langle z, w \rangle_n(t), \langle u, v \rangle_n(t_1) + \langle z, w \rangle_n(t)) \\ &\quad + d_\infty^n(\langle u, v \rangle_n(t_1), \langle u, v \rangle(t)) \\ &= d_\infty^n(\langle z, w \rangle_n(t), \langle z, w \rangle(t)) + d_\infty^n(\langle u, v \rangle_n(t_1), \langle u, v \rangle(t_1)) \\ &\quad + d_\infty^n(\langle u, v \rangle_n(t), \langle u, v \rangle(t)) \end{aligned}$$

$\forall t \in [t_1, t_1 + \delta(t_1)]$

Let  $n \rightarrow \infty$ . It follows that

$$\langle u, v \rangle(t_1) \oplus \langle z, w \rangle(t) \equiv \langle u, v \rangle(t) \quad \text{for all } t \in [t_1, t_1 + \delta(t_1)].$$

Hence the H-differences  $\langle u, v \rangle(t) \ominus \langle u, v \rangle(t_1)$  exist for all  $t_1 \in [t_1, t_1 + \delta(t_1)]$

Thus from (3.11) we have

$$d_\infty^n \left( \frac{\langle u, v \rangle(t) \ominus \langle u, v \rangle(t_1)}{t - t_1}, f(t_1, \langle u, v \rangle(t_1)) \right) \leq \varepsilon, \quad t_1 < t \leq t_1 + \delta(t_1)$$

So  $\lim_{t \rightarrow t_1^+} \frac{\langle u, v \rangle(t) \ominus \langle u, v \rangle(t_1)}{t - t_1} = f(t_1, \langle u, v \rangle(t_1))$ . Similarly, we have

$$\lim_{t \rightarrow t_1^-} \frac{\langle u, v \rangle(t) \ominus \langle u, v \rangle(t_1)}{t - t_1} = f(t_1, \langle u, v \rangle(t_1))$$

Hence  $\langle u, v \rangle'(t_1)$  exists and

$$\langle u, v \rangle'(t_1) = f(t_1, \langle u, v \rangle(t_1))$$

From  $t_1 \in [t_0, t_0 + r]$  is arbitrary, we know that Eq. (3.4) holds true and

$$\langle u, v \rangle \in C^1[[t_0, t_0 + r], B(\langle u, v \rangle_0, q)].$$

Thus, we conclude the proof. □

**Corollary 3.1.** *If we replace condition (3.2) by*

$$\begin{aligned} j\langle u, v \rangle'_{n+1}(t) &= jf(t, \langle u, v \rangle_n(t)) + B_n(t), \quad \langle u, v \rangle(t_0) = \langle u, v \rangle_0, \\ \| B_n(t) \| &\leq \varepsilon_n \quad \forall t \in [t_0, t_0 + r] (n = 0, 1, \dots) \end{aligned} \quad (3.12)$$

*and retain other assumptions, then the conclusions also hold true.*

*Proof 2.* This is completely similar to the proof of Theorem 3.1. □

## 4 Existence and uniqueness theorem for a solution

**Theorem 4.1.** *Let*

(a)  $f \in C[R_0, IF_n]$  and  $d_\infty^n \left( f(t, \langle u, v \rangle), 0_{(1,0)} \right) \leq M$  for all  $(t, \langle u, v \rangle) \in R_0$ .

(b)  $g \in C[[t_0, t_0 + p] \times [0, q], R]$ ,  $g(t, 0) \equiv 0$  and  $g(t, x) \leq M_1$ , for all  $t \in [t_0, t_0 + p]$ ,  $0 \leq x \leq q$  such that  $g(t, x)$  is nondecreasing on  $x$  (i.e.,  $t_0 \leq t \leq t_0 + p$ ,  $0 \leq x_1 \leq x_2 \leq q \implies g(t, x_1) \leq g(t, x_2)$ ), the initial value problem

$$x'(t) = g(t, x(t)), \quad x(t_0) = 0 \quad (4.1)$$

*has only the solution  $x(t) \equiv 0$  on  $[t_0, t_0 + p]$ .*

(c)  $d_\infty^n \left( f(t, \langle u, v \rangle), f(t, \langle u', v' \rangle) \right) \leq g \left( t, d_\infty^n \left( \langle u, v \rangle, \langle u', v' \rangle \right) \right)$ , for all  $(t, \langle u, v \rangle), (t, \langle u', v' \rangle) \in R_0$ , and  $d_\infty^n \left( \langle u, v \rangle, \langle u', v' \rangle \right) \leq q$ .

Then the Cauchy problem (3.4) has a unique solution  $\langle u, v \rangle \in C^1[[t_0, t_0 + r], B(x_0, q)]$  on  $[t_0, t_0 + r]$ , where  $r = \min\{p, q/M, q/M_1\}$ , and the successive iterations

$$\langle u, v \rangle_0(t) = \langle u, v \rangle_0, \quad \langle u, v \rangle_{n+1}(t) = \langle u, v \rangle_0 \oplus \int_{t_0}^t f(s, \langle u, v \rangle_n(s)) ds \quad (n = 0, 1, 2, \dots) \quad (4.2)$$

uniformly converge to  $\langle u, v \rangle(t)$  on  $[t_0, t_0 + r]$ .

*Proof 3.* from (4.2) and the assumption (a), by the inductive method we know

$$\begin{aligned} d_\infty^n \left( \langle u, v \rangle_{n+1}(t), \langle u, v \rangle_0 \right) &\leq \int_{t_0}^t d_\infty^n \left( f(s, \langle u, v \rangle_n(s)), 0_{(1,0)} \right) ds \\ &\leq q \quad \forall t \in [t_0, t_0 + r] \quad n = 0, 1, 2, \dots \end{aligned} \quad (4.3)$$

Hence  $\langle u, v \rangle_{n+1} \in C^1[[t_0, t_0 + r], B(x_0, q)]$  and

$$\langle u, v \rangle'_{n+1}(t) = f(t, (\langle u, v \rangle_n(t))), \quad \langle u, v \rangle_n(t_0) = \langle u, v \rangle_0 \quad (n = 0, 1, 2, \dots) \quad (4.4)$$

Let  $M_2 = \max\{M, M_1\}$ . Then  $r = \min\{p, q/M_2\}$ . and we get the successive iterations as

$$\begin{cases} x_0(t) = M_2(t - t_0) & t_0 \leq t \leq t_0 + r \\ x_{n+1}(t) = \int_{t_0}^t g(s, x_n(s)), & t_0 \leq t \leq t_0 + r \quad (n = 0, 1, 2, \dots) \end{cases} \quad (4.5)$$

It is immediate that

$$x_1(t) = \int_{t_0}^t g(s, x_0(s)) \leq M_1(t - t_0) \leq x_0(t) \leq q, \quad \forall t \in [t_0, t_0 + r] \quad (4.6)$$

So, by the inductive method and in view that  $g(t, x)$  is nondecreasing on  $x$ , we have

$$0 \leq x_{n+1}(t) \leq x_n(t) \leq q, \quad \forall t \in [t_0, t_0 + r] \quad (n = 0, 1, 2, \dots) \quad (4.7)$$

As  $|x'_{n+1}(t)| = |g(t, x_n(t))| \leq M_1$ , from the Ascoli-Arzelà theorem and (4.7) we know that  $\{x_n(t)\}$  uniformly converges to some continuous function  $x(t)$  on  $[t_0, t_0 + r]$  and

$$x(t) = \int_{t_0}^t g(s, x(s)) ds.$$

Thus  $x \in C^1[[t_0, t_0 + r], [0, q]]$  and  $x$  is the solution the initial value problem (4.1). From assumption (b) we get  $x(t) \equiv 0$ . In addition, we have

$$\begin{aligned} d_\infty^n \left( \langle u, v \rangle_1(t), \langle u, v \rangle_0 \right) &= d_\infty^n \left( \int_{t_0}^t f(s, \langle u, v \rangle_0(s)) ds, 0_{(1,0)} \right) \\ &\leq \int_{t_0}^t d_\infty^n \left( f(s, \langle u, v \rangle_0(s)), 0_{(1,0)} \right) ds \leq M(t - t_0) \leq x_0(t) \end{aligned}$$

Suppose  $d_\infty^n \left( \langle u, v \rangle_k(t), \langle u, v \rangle_{k-1} \right) \leq x_{k-1}(t)$ , then by the assumption (c), we have

$$\begin{aligned} d_\infty^n \left( \langle u, v \rangle_{k+1}(t), \langle u, v \rangle_k(t) \right) &= d_\infty^n \left( \int_{t_0}^t f(s, \langle u, v \rangle_k(s)) ds, \int_{t_0}^t f(s, \langle u, v \rangle_{k-1}(s)) ds \right) \\ &\leq \int_{t_0}^t d_\infty^n \left( f(s, \langle u, v \rangle_k(s)), f(s, \langle u, v \rangle_{k-1}(s)) \right) ds \leq \int_{t_0}^t g \left( s, d_\infty^n \left( \langle u, v \rangle_k(s), \langle u, v \rangle_{k-1}(s) \right) \right) ds \\ &\leq \int_{t_0}^t g \left( s, x_{k-1}(s) \right) ds = x_k(t) \end{aligned}$$



Thus by the inductive method we know

$$d_\infty^n \left( \langle u, v \rangle_{n+1}(t), \langle u, v \rangle_n(t) \right) \leq x_n(t) \quad t_0 \leq t \leq t_0 + r \quad (n = 0, 1, 2, \dots). \quad (4.8)$$

So, we have

$$\begin{aligned} d_\infty^n \left( \langle u, v \rangle'_{n+1}(t), \langle u, v \rangle'_n(t) \right) &= d_\infty^n \left( f(t, \langle u, v \rangle_n(t)), f(t, \langle u, v \rangle_{n-1}(t)) \right) \\ &\leq g \left( t, d_\infty^n \left( \langle u, v \rangle_n(t), \langle u, v \rangle_{n-1}(t) \right) \right) \\ &\leq g(t, x_{n-1}(t)) \end{aligned} \quad (4.9)$$

Assume  $m \geq n$ , and in view of (4.9) and (4.7) we get

$$\begin{aligned} d_\infty^n \left( \langle u, v \rangle'_n(t), \langle u, v \rangle'_m(t) \right) &\leq d_\infty^n \left( f(t, \langle u, v \rangle_{n-1}(t)), f(t, \langle u, v \rangle_n(t)) \right) \\ &\quad + d_\infty^n \left( f(t, \langle u, v \rangle_n(t)), f(t, \langle u, v \rangle_m(t)) \right) \\ &\quad + d_\infty^n \left( f(t, \langle u, v \rangle_m(t)), f(t, \langle u, v \rangle_{m-1}(t)) \right) \\ &\leq g(t, x_{n-1}(t)) + g(t, d_\infty^n(\langle u, v \rangle_n(t), \langle u, v \rangle_m(t))) + g(t, x_{m-1}(t)) \\ &\leq 2g(t, x_{n-1}(t)) + g(t, d_\infty^n(\langle u, v \rangle_n(t), \langle u, v \rangle_m(t))) \end{aligned}$$

Furthermore, from

$$\begin{aligned} &d_\infty^n \left( \langle u, v \rangle_n(t+h), \langle u, v \rangle_m(t+h) \right) \\ &\leq d_\infty^n \left( \langle u, v \rangle_n(t+h), \langle u, v \rangle_m(t+h) - \langle u, v \rangle_m(t) + \langle u, v \rangle_n(t) \right) \\ &\quad + d_\infty^n \left( \langle u, v \rangle_m(t+h) \ominus \langle u, v \rangle_m(t) + \langle u, v \rangle_n(t), \langle u, v \rangle_m(t+h) \right) \\ &= d_\infty^n \left( \langle u, v \rangle_n(t+h) \ominus \langle u, v \rangle_n(t), \langle u, v \rangle_m(t+h) \ominus \langle u, v \rangle_m(t) \right) \\ &\quad + d_\infty^n \left( \langle u, v \rangle_n(t), \langle u, v \rangle_m(t) \right) \end{aligned}$$

we deduce that

$$\begin{aligned} &D^+ d_\infty^n \left( \langle u, v \rangle_n(t), \langle u, v \rangle_m(t) \right) \\ &= \lim_{h \rightarrow 0^+} \frac{d_\infty^n(\langle u, v \rangle_n(t+h), \langle u, v \rangle_m(t+h)) - d_\infty^n(\langle u, v \rangle_n(t), \langle u, v \rangle_m(t))}{h} \\ &\leq \lim_{h \rightarrow 0^+} \frac{d_\infty^n(\langle u, v \rangle_n(t+h) \ominus \langle u, v \rangle_n(t), \langle u, v \rangle_m(t+h) \ominus \langle u, v \rangle_m(t))}{h} \\ &= d_\infty^n \left( \langle u, v \rangle'_n(t), \langle u, v \rangle'_m(t) \right) < 2g(t, x_{n-1}(t)) + g(t, d_\infty^n(\langle u, v \rangle_n(t), \langle u, v \rangle_m(t))) \end{aligned}$$

Since  $g(t, x_{n-1}(t))$  uniformly converges to 0, then for arbitrary  $\varepsilon > 0$  there exists a natural number  $N$  such that

$$D^+ d_\infty^n \left( \langle u, v \rangle_n(t), \langle u, v \rangle_m(t) \right) < g(t, d_\infty^n(\langle u, v \rangle_n(t), \langle u, v \rangle_m(t))) + \varepsilon \quad \forall m \geq n > N$$

Here  $D^+$  is the Dini derivative (see [7]). From the fact that  $d_\infty^n \left( \langle u, v \rangle_n(t_0), \langle u, v \rangle_m(t_0) \right) = 0 < \varepsilon$  and by proposition 2.1, we have

$$d_\infty^n \left( \langle u, v \rangle_n(t), \langle u, v \rangle_m(t) \right) \leq w(t, \varepsilon) \quad \forall t \in [t_0, t_0 + r] \quad \forall m \geq n > N \quad (4.10)$$

where  $w(t, \varepsilon)$  is the maximum solution to the initial value problem

$$x'(t) = g(t, x(t)) + \varepsilon, \quad x(t_0) = \varepsilon \quad (4.11)$$

By proposition 2.2 we know that  $w(t, \varepsilon)$  uniformly converges to the maximum solution  $x(t) \equiv 0$  of problem (4.1) on  $t_0 \leq t \leq t_0 + r$  as  $\varepsilon \rightarrow 0$ .

Thus, according to (4.10) and that  $(IF_n, d_\infty^n)$  is complete, we know that there exists an intuitionistic fuzzy set-valued mapping  $\langle u, v \rangle : T \rightarrow IF_n$  such that  $d_\infty^n \left( \langle u, v \rangle_n(t), \langle u, v \rangle(t) \right)$  uniformly converges to 0 as  $n \rightarrow \infty$ . Applying (4.4) and Corollary (3.1) we have  $\langle u, v \rangle \in C^1[[t_0, t_0 + r], B(\langle u, v \rangle_0, q)]$  and  $\langle u, v \rangle(t)$  is the solution of the initial value problem (3.4).

Finally, we prove the uniqueness. Suppose  $\langle z, w \rangle(t)$  is another solution of initial value problem (3.4). Let

$$m(t) = d_\infty^n \left( \langle u, v \rangle(t), \langle z, w \rangle(t) \right)$$

Then  $m(t_0) \equiv 0$

$$D^+ m(t) \leq d_\infty^n \left( \langle u, v \rangle'(t), \langle z, w \rangle'(t) \right) = d_\infty^n \left( f(t, \langle u, v \rangle(t)), f(t, \langle z, w \rangle(t)) \right) \leq g(t, m(t)).$$

Hence from proposition 2.2 we know

$$d_\infty^n \left( \langle u, v \rangle(t), \langle z, w \rangle(t) \right) \leq x(t) \equiv 0, \quad \forall t \in [t_0, t_0 + r]$$

where  $x(t) \equiv 0$  is the maximum solution of problem (4.1) on  $[t_0, t_0 + r]$ .

Therefore  $\langle u, v \rangle(t) = \langle z, w \rangle(t)$ . □

**Corollary 4.1.** *Let  $f \in C[R_0, IF_n]$  such that  $d_\infty^n \left( f(t, \langle u, v \rangle), 0 \right) \leq M$  for all  $(t, \langle u, v \rangle) \in R_0$  and  $f$  satisfies the Lipschitz condition*

$$d_\infty^n \left( f(t, \langle u, v \rangle), f(t, \langle u', v' \rangle) \right) \leq L d_\infty^n \left( \langle u, v \rangle, \langle u', v' \rangle \right), \quad \forall (t, \langle u, v \rangle), (t, \langle u', v' \rangle) \in R_0$$

where  $L$  is a constant. Then the Cauchy problem (3.4) has an unique solution  $\langle u, v \rangle \in C^1[[t_0, t_0 + r], B(x_0, q)]$  on  $[t_0, t_0 + r]$ , where  $r = \min\{p, q/M, 1/L\}$ , and the successive iterations (4.2) uniformly converge to  $\langle u, v \rangle(t)$  on  $[t_0, t_0 + r]$ .

*Proof 4.* In the proof of Theorem (4.1), taking  $g(t, x) = L.x$  we then obtain the proof of Corollary 4.1, where  $M_1 = L.q$ , hence  $r = \min\{p, q/M, 1/L\}$ . □

## References

- [1] Atanassov, K. (1983) Intuitionistic Fuzzy Sets, *VII ITKR Session*, Sofia, 20-23 June 1983 (Deposited in Centr. Sci.-Techn. Library of the Bulg. Acad. of Sci., 1697/84) (in Bulgarian). Reprinted: *Int. J. Bioautomation*, 2016, 20(S1), S1–S6.

- [2] Atanassov, K. (1986) Intuitionistic fuzzy sets, *Fuzzy Sets and Systems*, 20(1), 87–96.
- [3] Atanassov, K. (1994) Operators over interval valued intuitionistic fuzzy sets, *Fuzzy Sets and Systems*, 64(2), 159–174.
- [4] De, S. K., Biswas, R., & Roy, A. R. (2001) An application of intuitionistic fuzzy sets in medical diagnosis, *Fuzzy Sets and Systems*, 117(2), 209–213.
- [5] El Allaoui, A., Melliani, S., & Chadli, L.S. (2016) The Cauchy problem for complex intuitionistic fuzzy differential equations, *Notes on Intuitionistic Fuzzy Sets*, 22(4), 53–63.
- [6] Kharal, A. (2009) Homeopathic drug selection using intuitionistic fuzzy sets, *Homeopathy*, 98(1), 35–39.
- [7] Lakshmikantham, V., & Leela, S. (1969) *Differential and Integral Inequalities*, Vols. I and II, Academic Press, New York.
- [8] Li, D.F. & Cheng, C. T. (2002) New similarity measures of intuitionistic fuzzy sets and application to pattern recognitions, *Pattern Recognit. Lett.*, 23(1–3), 221–225.
- [9] Li, D. F. (2005) Multiattribute decision making models and methods using intuitionistic fuzzy sets, *J. Comput. Syst. Sci.*, 70, 73–85.
- [10] Klement, E. P., Puri, M. L., & Ralescu, D. A. (1986) Limit Theorems for Fuzzy Random Variables, *Proc. R. Soc. Lond. A*, 407, 171–182.
- [11] Melliani, S., & Chadli, L. S. (2000) Intuitionistic fuzzy differential equation. Part 1, *Notes on Intuitionistic Fuzzy Sets*, 6(2), 37–41.
- [12] Melliani, S., Elomari, M., Chadli, L. S., & Ettoussi, R. (2015) Intuitionistic Fuzzy Metric Space, *Notes on Intuitionistic Fuzzy sets*, 21(1), 43–53.
- [13] Melliani, S., Elomari, M., Chadli, L. S., & Ettoussi, R. (2015) Intuitionistic fuzzy fractional equation, *Notes on Intuitionistic Fuzzy Sets*, 21(4), 76–89.
- [14] Wu, C., & Song, S. (1996) Approximate Solutions, Existence, and Uniqueness of the Cauchy Problem of Fuzzy Differential Equations, *Journal Of Mathematical Analysis And Applications*, 202, 629–644.
- [15] Zadeh, L. A. (1965) Fuzzy sets, *Inf. Control*, 8, 338–353.