

Remark on Dworniczak’s intuitionistic fuzzy implications. Part 1

Lilija Atanassova

Institute of Information and Communication Technologies

Bulgarian Academy of Sciences,

Acad. G. Bonchev Str., Bl. 2, Sofia–1113, Bulgaria

e-mail: l.c.atanassova@gmail.com

Abstract: Four new intuitionistic fuzzy implications are introduced. They are modal forms of Dworniczak’s intuitionistic fuzzy implication $\rightarrow_{150,\lambda}$. Their basic properties are studied.

Keywords: Implication, Intuitionistic fuzzy logic

AMS Classification: Primary 03E72; Secondary 04E72.

1 Introduction

In [7, 8, 9], Piotr Dworniczak introduce three intuitionistic fuzzy implications, that generalized the defined by the author intuitionistic fuzzy implications in [4, 5, 6]. Some of their properties are discussed in [1].

Here, following the idea from [2], we introduce four new intuitionistic fuzzy implications, modifying Dworniczak’s intuitionistic fuzzy implication

$$\langle a, b \rangle \rightarrow_{150,\lambda} \langle c, d \rangle = \left\langle \frac{b + c + \lambda - 1}{2\lambda}, \frac{a + d + \lambda - 1}{2\lambda} \right\rangle,$$

where $\lambda \geq 1$.

2 Main results

Let everywhere below the variables x and y have truth values $\langle a, b \rangle$ and $\langle C, D \rangle$, where $a, b, c, d \in [0, 1]$, $a + b \leq 1$, $c + d \leq 1$.

In [2, 3] the following operations and operators are defined:

$$\neg \langle a, b \rangle \equiv \neg_1 \langle a, b \rangle = \langle b, a \rangle,$$

$$\langle a, b \rangle @ \langle c, d \rangle = \left\langle \frac{a+c}{2}, \frac{b+d}{2} \right\rangle,$$

$$\square \langle a, b \rangle = \langle a, 1-a \rangle,$$

$$\diamond \langle a, b \rangle = \langle 1-b, b \rangle,$$

$$\boxplus_{\alpha, \beta, \gamma} \langle a, b \rangle = \langle \alpha a, \beta b + \gamma \rangle,$$

$$\boxtimes_{\alpha, \beta, \gamma} \langle a, b \rangle = \langle \alpha a + \gamma, \beta b \rangle,$$

where $\alpha, \beta, \gamma \in [0, 1]$ and $\max(\alpha, \beta) + \gamma \leq 1$,

$$\blacksquare_{\alpha, \beta, \gamma, \delta} \langle a, b \rangle = \langle \alpha a + \gamma, \beta b + \delta \rangle,$$

where $\alpha, \beta, \gamma, \delta \in [0, 1]$ and $\max(\alpha, \beta) + \gamma + \delta \leq 1$.

We use the following formulas:

$$\langle a, b \rangle \rightarrow_{150, \lambda}^1 \langle c, d \rangle = \square \langle a, b \rangle \rightarrow_{150, \lambda} \square \langle c, d \rangle,$$

$$\langle a, b \rangle \rightarrow_{150, \lambda}^2 \langle c, d \rangle = \square \langle a, b \rangle \rightarrow_{150, \lambda} \diamond \langle c, d \rangle,$$

$$\langle a, b \rangle \rightarrow_{150, \lambda}^3 \langle c, d \rangle = \diamond \langle a, b \rangle \rightarrow_{150, \lambda} \diamond \langle c, d \rangle,$$

$$\langle a, b \rangle \rightarrow_{150, \lambda}^4 \langle c, d \rangle = \diamond \langle a, b \rangle \rightarrow_{150, \lambda} \square \langle c, d \rangle.$$

So, we obtain the explicit forms of the new four implications as follows:

$$\langle a, b \rangle \rightarrow_{150, \lambda}^1 \langle c, d \rangle = \left\langle \frac{-a+c+\lambda}{2\lambda}, \frac{a-c+\lambda}{2\lambda} \right\rangle,$$

$$\langle a, b \rangle \rightarrow_{150, \lambda}^2 \langle c, d \rangle = \left\langle \frac{1-a-d+\lambda}{2\lambda}, \frac{a+d+\lambda-1}{2\lambda} \right\rangle,$$

$$\langle a, b \rangle \rightarrow_{150, \lambda}^3 \langle c, d \rangle = \left\langle \frac{b+c+\lambda-1}{2\lambda}, \frac{1-b-c+\lambda}{2\lambda} \right\rangle,$$

$$\langle a, b \rangle \rightarrow_{150, \lambda}^4 \langle c, d \rangle = \left\langle \frac{b-d+\lambda}{2\lambda}, \frac{-b+d+\lambda}{2\lambda} \right\rangle.$$

First, we check that for every $i = 1, 2, 3, 4$ and for every $\lambda \geq 1$:

$$\langle 0, 1 \rangle \rightarrow_{150, \lambda}^i \langle 0, 1 \rangle = \left\langle \frac{1}{2}, \frac{1}{2} \right\rangle,$$

$$\langle 0, 1 \rangle \rightarrow_{150, \lambda}^i \langle 1, 0 \rangle = \left\langle \frac{\lambda+1}{2}, \frac{\lambda-1}{2} \right\rangle,$$

$$\langle 1, 0 \rangle \rightarrow_{150, \lambda}^i \langle 0, 1 \rangle = \left\langle \frac{\lambda-1}{2}, \frac{\lambda+1}{2} \right\rangle,$$

$$\langle 1, 0 \rangle \rightarrow_{150, \lambda}^i \langle 1, 0 \rangle = \left\langle \frac{1}{2}, \frac{1}{2} \right\rangle.$$

Second, we check that for every $\lambda \geq 1$:

$$\langle 0, 0 \rangle \rightarrow_{150, \lambda}^1 \langle 0, 1 \rangle = \left\langle \frac{1}{2}, \frac{1}{2} \right\rangle,$$

$$\begin{aligned}
\langle 0, 0 \rangle \rightarrow_{150, \lambda}^2 \langle 0, 1 \rangle &= \langle \frac{1}{2}, \frac{1}{2} \rangle, \\
\langle 0, 0 \rangle \rightarrow_{150, \lambda}^3 \langle 0, 1 \rangle &= \langle \frac{\lambda - 1}{2}, \frac{\lambda + 1}{2} \rangle, \\
\langle 0, 0 \rangle \rightarrow_{150, \lambda}^4 \langle 0, 1 \rangle &= \langle \frac{\lambda - 1}{2}, \frac{\lambda + 1}{2} \rangle, \\
\langle 0, 0 \rangle \rightarrow_{150, \lambda}^1 \langle 0, 0 \rangle &= \langle \frac{1}{2}, \frac{1}{2} \rangle, \\
\langle 0, 0 \rangle \rightarrow_{150, \lambda}^2 \langle 0, 0 \rangle &= \langle \frac{\lambda + 1}{2}, \frac{\lambda - 1}{2} \rangle, \\
\langle 0, 0 \rangle \rightarrow_{150, \lambda}^3 \langle 0, 0 \rangle &= \langle \frac{\lambda - 1}{2}, \frac{\lambda + 1}{2} \rangle, \\
\langle 0, 0 \rangle \rightarrow_{150, \lambda}^4 \langle 0, 0 \rangle &= \langle \frac{1}{2}, \frac{1}{2} \rangle, \\
\langle 0, 0 \rangle \rightarrow_{150, \lambda}^1 \langle 1, 0 \rangle &= \langle \frac{\lambda + 1}{2}, \frac{\lambda - 1}{2} \rangle, \\
\langle 0, 0 \rangle \rightarrow_{150, \lambda}^2 \langle 1, 0 \rangle &= \langle \frac{\lambda + 1}{2}, \frac{\lambda - 1}{2} \rangle, \\
\langle 0, 0 \rangle \rightarrow_{150, \lambda}^3 \langle 1, 0 \rangle &= \langle \frac{1}{2}, \frac{1}{2} \rangle, \\
\langle 0, 0 \rangle \rightarrow_{150, \lambda}^4 \langle 1, 0 \rangle &= \langle \frac{1}{2}, \frac{1}{2} \rangle, \\
\langle 0, 1 \rangle \rightarrow_{150, \lambda}^1 \langle 0, 0 \rangle &= \langle \frac{1}{2}, \frac{1}{2} \rangle, \\
\langle 0, 1 \rangle \rightarrow_{150, \lambda}^2 \langle 0, 0 \rangle &= \langle \frac{\lambda + 1}{2}, \frac{\lambda - 1}{2} \rangle, \\
\langle 0, 1 \rangle \rightarrow_{150, \lambda}^3 \langle 0, 0 \rangle &= \langle \frac{1}{2}, \frac{1}{2} \rangle, \\
\langle 0, 1 \rangle \rightarrow_{150, \lambda}^4 \langle 0, 0 \rangle &= \langle \frac{\lambda + 1}{2}, \frac{\lambda - 1}{2} \rangle, \\
\langle 1, 0 \rangle \rightarrow_{150, \lambda}^1 \langle 0, 0 \rangle &= \langle \frac{\lambda - 1}{2}, \frac{\lambda + 1}{2} \rangle, \\
\langle 1, 0 \rangle \rightarrow_{150, \lambda}^2 \langle 0, 0 \rangle &= \langle \frac{1}{2}, \frac{1}{2} \rangle, \\
\langle 1, 0 \rangle \rightarrow_{150, \lambda}^3 \langle 0, 0 \rangle &= \langle \frac{\lambda - 1}{2}, \frac{\lambda + 1}{2} \rangle, \\
\langle 1, 0 \rangle \rightarrow_{150, \lambda}^4 \langle 0, 0 \rangle &= \langle \frac{1}{2}, \frac{1}{2} \rangle.
\end{aligned}$$

Using definition from [2, 3]

$$\langle a, b \rangle \geq \langle c, d \rangle \text{ if and only if } a \geq c \text{ and } b \leq d,$$

we can prove the validity of the following theorem.

Theorem 1. For every $a, b, c, d \in [0, 1]$, so that $a + b \leq 1$ and $c + d \leq 1$ and for every $\lambda \geq 1$:

$$\langle a, b \rangle \rightarrow_{150, \lambda}^2 \langle c, d \rangle \geq \langle a, b \rangle \rightarrow_{150, \lambda}^1 \langle c, d \rangle, \quad (1)$$

$$\langle a, b \rangle \rightarrow_{150, \lambda}^4 \langle c, d \rangle \geq \langle a, b \rangle \rightarrow_{150, \lambda}^3 \langle c, d \rangle, \quad (2)$$

$$\langle a, b \rangle \rightarrow_{150, \lambda}^2 \langle c, d \rangle \geq \langle a, b \rangle \rightarrow_{150, \lambda}^4 \langle c, d \rangle, \quad (3)$$

$$\langle a, b \rangle \rightarrow_{150, \lambda}^1 \langle c, d \rangle \geq \langle a, b \rangle \rightarrow_{150, \lambda}^3 \langle c, d \rangle. \quad (4)$$

Proof: For example, let us check the validity of the fourth inequality.

First, we see, that

$$-a + c + \lambda - b - c - \lambda + 1 = 1 - a - b \geq 0,$$

$$1 - b - c + \lambda - a + c - \lambda = 1 - a - b \geq 0.$$

Therefore,


$$\left\langle \frac{-a + c + \lambda}{2\lambda}, \frac{a - c + \lambda}{2\lambda} \right\rangle \geq \left\langle \frac{b + c + \lambda - 1}{2\lambda}, \frac{1 - b - c + \lambda}{2\lambda} \right\rangle,$$

i.e.,

$$\langle a, b \rangle \rightarrow_{150, \lambda}^1 \langle c, d \rangle \geq \langle a, b \rangle \rightarrow_{150, \lambda}^3 \langle c, d \rangle.$$

Hence, (4) is valid. (1) – (3) are proved by analogy. \square

Now, we can construct the following diagram



Now, we show that the new implications can be represented by a part of the above operators.

Theorem 2. For every $a, b, c, d \in [0, 1]$, so that $a + b \leq 1$ and $c + d \leq 1$ and for every $\lambda \geq 1$:

$$\langle a, b \rangle \rightarrow_{150, \lambda}^1 \langle c, d \rangle = \boxtimes_{\frac{1}{\lambda}, \frac{1}{\lambda}, \frac{\lambda-1}{\lambda}} \diamond \neg \langle a, b \rangle @ \boxplus_{\frac{1}{\lambda}, \frac{1}{\lambda}, \frac{\lambda-1}{\lambda}} \square \langle c, d \rangle,$$

$$\langle a, b \rangle \rightarrow_{150, \lambda}^2 \langle c, d \rangle = \boxtimes_{\frac{1}{\lambda}, \frac{1}{\lambda}, \frac{\lambda-1}{\lambda}} \diamond \neg \langle a, b \rangle @ \boxplus_{\frac{1}{\lambda}, \frac{1}{\lambda}, \frac{\lambda-1}{\lambda}} \square \neg \langle c, d \rangle,$$

$$\langle a, b \rangle \rightarrow_{150, \lambda}^1 \langle c, d \rangle = \boxplus_{\frac{1}{\lambda}, \frac{1}{\lambda}, \frac{\lambda-1}{\lambda}} \square \neg \langle a, b \rangle @ \boxtimes_{\frac{1}{\lambda}, \frac{1}{\lambda}, \frac{\lambda-1}{\lambda}} \diamond \langle c, d \rangle,$$

$$\langle a, b \rangle \rightarrow_{150, \lambda}^3 \langle c, d \rangle = \boxtimes_{\frac{1}{\lambda}, \frac{1}{\lambda}, \frac{\lambda-1}{\lambda}} \square \neg \langle a, b \rangle @ \boxplus_{\frac{1}{\lambda}, \frac{1}{\lambda}, \frac{\lambda-1}{\lambda}} \diamond \langle c, d \rangle.$$

The proofs of these assertions are similar to the above one. □

Corollary. For every $a, b, c, d \in [0, 1]$, so that $a + b \leq 1$ and $c + d \leq 1$ and for every $\lambda \geq 1$:

$$\langle a, b \rangle \rightarrow_{150, \lambda}^1 \langle c, d \rangle = \blacksquare_{\frac{1}{\lambda}, \frac{1}{\lambda}, \frac{\lambda-1}{\lambda}, 0} \diamond \neg \langle a, b \rangle @ \bullet_{\frac{1}{\lambda}, \frac{1}{\lambda}, 0, \frac{\lambda-1}{\lambda}} \square \langle c, d \rangle,$$

$$\langle a, b \rangle \rightarrow_{150, \lambda}^2 \langle c, d \rangle = \bullet_{\frac{1}{\lambda}, \frac{1}{\lambda}, \frac{\lambda-1}{\lambda}, 0} \diamond \neg \langle a, b \rangle @ \blacksquare_{\frac{1}{\lambda}, \frac{1}{\lambda}, 0, \frac{\lambda-1}{\lambda}} \square \langle c, d \rangle,$$

$$\langle a, b \rangle \rightarrow_{150, \lambda}^1 \langle c, d \rangle = \blacksquare_{\frac{1}{\lambda}, \frac{1}{\lambda}, 0, \frac{\lambda-1}{\lambda}} \square \neg \langle a, b \rangle @ \bullet_{\frac{1}{\lambda}, \frac{1}{\lambda}, \frac{\lambda-1}{\lambda}, 0} \diamond \langle c, d \rangle,$$

$$\langle a, b \rangle \rightarrow_{150, \lambda}^3 \langle c, d \rangle = \bullet_{\frac{1}{\lambda}, \frac{1}{\lambda}, 0, \frac{\lambda-1}{\lambda}} \square \neg \langle a, b \rangle @ \blacksquare_{\frac{1}{\lambda}, \frac{1}{\lambda}, \frac{\lambda-1}{\lambda}, 0} \diamond \langle c, d \rangle.$$

3 Conclusion

In the next parts of the research, we will study from one side other properties of the new implications and from another – the modifications of the two other Dworniczak's implications and their properties.

References

- [1] Angelova, N., Marinov, E., & Atanassov, K. (2015) Intuitionistic Fuzzy Implications and Kolmogorov's and Lukasiewicz-Tarski's Axioms of Logic. *Notes on Intuitionistic Fuzzy Sets*, 21(2), 35–42.
- [2] Atanassov, K. (2012) *On Intuitionistic Fuzzy Sets Theory*, Springer, Berlin.
- [3] Atanassov, K., Szmidt E., & Kacprzyk J. (2013) On intuitionistic fuzzy pairs. *Notes on Intuitionistic Fuzzy Sets*, 19(3), 1–13.
- [4] Atanassova, L. (2009) A new intuitionistic fuzzy implication, *Cybernetics and Information Technologies*, 9(2), 21–25.
- [5] Atanassova, L. (2012) On two modifications of the intuitionistic fuzzy implication $\rightarrow_{@}$, *Notes on Intuitionistic Fuzzy Sets*, 18(2), 26–30.
- [6] Atanassova, L., On the modal form of the intuitionistic fuzzy implications $\rightarrow'_{@}$ and $\rightarrow''_{@}$, *Issues in Intuitionistic Fuzzy Sets and Generalized Nets*, 10, 5–11.

- [7] Dworniczak, P. (2010) Some remarks about the L. Atanassova's paper "A new intuitionistic fuzzy implication", *Cybernetics and Information Technologies*, 10(3), 3–9.
- [8] Dworniczak, P. (2010) On one class of intuitionistic fuzzy implications, *Cybernetics and Information Technologies*, 10(4), 13–21.
- [9] Dworniczak, P. (2011) On some two-parametric intuitionistic fuzzy implication, *Notes on Intuitionistic Fuzzy Sets*, 17(2), 8–16.