

## Remark on the intuitionistic fuzzy forms of two classical logic axioms. Part 2

Lilija Atanassova

Institute of Information and Communication Technologies  
Bulgarian Academy of Sciences  
Acad. G. Bonchev Str., Bl. 2, Sofia-1113, Bulgaria  
e-mail: l.c.atanassova@gmail.com

**Abstract:** .....

**Keywords:** Intuitionistic fuzzy implication, Logical tautology.

**AMS Classification:** 03E72.

### 1 Introduction

In a series of research, we discuss the intuitionistic fuzzy forms of some classical logic axioms, and check their validity in the case of intuitionistic fuzziness.

In [?] we determined the implications that satisfy the standard logical tautologies

$$(p \& q) \rightarrow r = (p \rightarrow (q \rightarrow r)),$$

$$p \rightarrow q = (p \rightarrow (p \rightarrow q)).$$

Two other well-known logical tautologies (see, e.g., [?]) are

$$(p \vee q) \rightarrow r = (p \rightarrow r) \& (q \rightarrow r), \quad (1)$$

$$(p \& q) \rightarrow r = (p \rightarrow r) \vee (q \rightarrow r). \quad (2)$$

Here, we discuss their validity for the different cases of intuitionistic fuzzy implications. In [?], 138 of them are given.

Below, we determine which of these 138 intuitionistic fuzzy implications satisfy (1) and (2).

**Theorem 1.** Implications  $\rightarrow_2, \rightarrow_3, \rightarrow_4, \rightarrow_5, \rightarrow_{12}, \rightarrow_{13}, \rightarrow_{14}, \rightarrow_{16}, \rightarrow_{18}, \rightarrow_{19}, \rightarrow_{20}, \rightarrow_{22}, \rightarrow_{23}, \rightarrow_{25}$

,  $\rightarrow_{26}, \rightarrow_{27}, \rightarrow_{28}, \rightarrow_{29}, \rightarrow_{31}, \rightarrow_{32}, \rightarrow_{33}, \rightarrow_{34}, \rightarrow_{35}, \rightarrow_{37}, \rightarrow_{40}, \rightarrow_{41}, \rightarrow_{42}, \rightarrow_{43}, \rightarrow_{44}, \rightarrow_{45},$   
 $\rightarrow_{47}, \rightarrow_{48}, \rightarrow_{49}, \rightarrow_{50}, \rightarrow_{52}, \rightarrow_{55}, \rightarrow_{56}, \rightarrow_{57}, \rightarrow_{58}, \rightarrow_{59}, \rightarrow_{60}, \rightarrow_{61}, \rightarrow_{64}, \rightarrow_{66}, \rightarrow_{67}, \rightarrow_{69}, \rightarrow_{70},$   
 $\rightarrow_{71}, \rightarrow_{72}, \rightarrow_{73}, \rightarrow_{74}, \rightarrow_{76}, \rightarrow_{77}, \rightarrow_{78}, \rightarrow_{79}, \rightarrow_{80}, \rightarrow_{81}, \rightarrow_{82}, \rightarrow_{83}, \rightarrow_{84}, \rightarrow_{85}, \rightarrow_{86}, \rightarrow_{87}, \rightarrow_{88},$   
 $\rightarrow_{89}, \rightarrow_{90}, \rightarrow_{91}, \rightarrow_{92}, \rightarrow_{93}, \rightarrow_{94}, \rightarrow_{95}, \rightarrow_{96}, \rightarrow_{97}, \rightarrow_{98}, \rightarrow_{99}, \rightarrow_{102}, \rightarrow_{105}, \rightarrow_{108}, \rightarrow_{124}, \rightarrow_{125},$   
 $\rightarrow_{127}, \rightarrow_{129}, \rightarrow_{130}, \rightarrow_{132}, \rightarrow_{134}, \rightarrow_{135}, \rightarrow_{137}$  satisfy (1).

**Proof.** Below, we prove that (1) is valid for implication  $\rightarrow_3$ . The rest assertions are proved by analogy. Let everywhere below, truth-values of  $p, q, r$  be:

$$V(p) = \langle a, b \rangle,$$

$$V(q) = \langle c, d \rangle,$$

$$V(r) = \langle e, f \rangle.$$

In [?], implication  $\rightarrow_3$  is Gödel's implication, that has the form:

$$V(p \rightarrow_3 q) = \langle a, b \rangle \rightarrow_3 \langle c, d \rangle = \langle 1 - (1 - c) \cdot \text{sg}(a - c), d \cdot \text{sg}(a - c) \rangle,$$

where

$$\text{sg}(x) = \begin{cases} 1, & \text{if } x > 0 \\ 0, & \text{if } x \leq 0 \end{cases}.$$

There it is defined that

$$V(p \& q) = \langle \min(a, c), \max(b, d) \rangle,$$

$$V(p \vee q) = \langle \max(a, c), \min(b, d) \rangle.$$

Therefore, the expression (1) has the form

$$(p \vee q) \rightarrow r = (p \rightarrow r) \& (q \rightarrow r)$$

or

$$(\langle a, b \rangle \vee \langle c, d \rangle) \rightarrow_3 \langle e, f \rangle = (\langle a, b \rangle \rightarrow_3 \langle e, f \rangle) \& (\langle c, d \rangle \rightarrow_3 \langle e, f \rangle).$$

The left side has the form

$$\begin{aligned} & (\langle a, b \rangle \vee \langle c, d \rangle) \rightarrow_3 \langle e, f \rangle \\ &= \langle \max(a, c), \min(b, d) \rangle \rightarrow_3 \langle e, f \rangle \\ &= \langle 1 - (1 - e) \cdot \text{sg}(\max(a, c) - e), f \cdot \text{sg}(\max(a, c) - e) \rangle. \end{aligned}$$

The right side has the form

$$\begin{aligned} & (\langle a, b \rangle \rightarrow_3 \langle e, f \rangle) \& (\langle c, d \rangle \rightarrow_3 \langle e, f \rangle) \\ &= \langle 1 - (1 - e) \cdot \text{sg}(a - e), f \cdot \text{sg}(a - e) \rangle \& \langle 1 - (1 - e) \cdot \text{sg}(c - e), f \cdot \text{sg}(c - e) \rangle \\ &= \langle \min(1 - (1 - e) \cdot \text{sg}(a - e), 1 - (1 - e) \cdot \text{sg}(c - e)), \max(f \cdot \text{sg}(a - e), f \cdot \text{sg}(c - e)) \rangle \end{aligned}$$

$$\begin{aligned}
&= \langle 1 - \max((1 - e) \cdot \text{sg}(a - e), (1 - e) \cdot \text{sg}(c - e)), f \cdot \max(\text{sg}(a - e), \text{sg}(c - e)) \rangle \\
&= \langle 1 - (1 - e) \cdot \max(\text{sg}(a - e), \text{sg}(c - e)), f \cdot \max(\text{sg}(a - e), \text{sg}(c - e)) \rangle.
\end{aligned}$$

Let

$$X \equiv (1 - (1 - e) \cdot \text{sg}(\max(a, c) - e)) - (1 - (1 - e) \cdot \max(\text{sg}(a - e), \text{sg}(c - e))).$$

Then

$$\begin{aligned}
X &= (1 - e) \cdot \max(\text{sg}(a - e), \text{sg}(c - e)) - (1 - e) \cdot \text{sg}(\max(a, c) - e) \\
&= (1 - e) \cdot (\max(\text{sg}(a - e), \text{sg}(c - e)) - \text{sg}(\max(a, c) - e)).
\end{aligned}$$

If  $a \leq c$ , then

$$X = (1 - e) \cdot (\max(\text{sg}(a - e), \text{sg}(c - e)) - \text{sg}(c - e)) \geq 0.$$

If  $a > e$ , then

$$X = (1 - e) \cdot (\max(1, 1) - 1) = 0.$$

If  $a \leq e < c$ , then

$$X = (1 - e) \cdot (\max(0, 1) - \text{sg}(\max(c - e))) = (1 - e) \cdot (1 - 1) = 0.$$

If  $c \leq e$ , then

$$X = (1 - e) \cdot (\max(0, 0) - \text{sg}(c - e)) = (1 - e) \cdot (0 - 0) = 0.$$

If  $a > c$ , then

$$X = (1 - e) \cdot (\max(\text{sg}(a - e), \text{sg}(c - e)) - \text{sg}(a - e)).$$

If  $a \leq e$ , then

$$X = (1 - e) \cdot (\max(0, 0) - 0) = 0.$$

If  $a > e \geq c$ , then

$$X = (1 - e) \cdot (\max(1, 0) - 1) = 0.$$

If  $a > c > e$ , then

$$X = (1 - e) \cdot (\max(1, 1) - 1) = 0.$$

Therefore, in all cases  $X = 0$ .

Analogously, we check that

$$Y \equiv f \cdot \text{sg}(\max(a, c) - e) - f \cdot \max(\text{sg}(a - e), \text{sg}(c - e)) = 0.$$

Hence, (1) is an equality for implication  $\rightarrow_3$ .

**Theorem 2.** Implications  $\rightarrow_2, \rightarrow_3, \rightarrow_4, \rightarrow_5, \rightarrow_8, \rightarrow_{11}, \rightarrow_{12}, \rightarrow_{13}, \rightarrow_{16}, \rightarrow_{18}, \rightarrow_{19}, \rightarrow_{20}, \rightarrow_{22}, \rightarrow_{23}, \rightarrow_{25}, \rightarrow_{26}, \rightarrow_{27}, \rightarrow_{28}, \rightarrow_{29}, \rightarrow_{31}, \rightarrow_{32}, \rightarrow_{33}, \rightarrow_{34}, \rightarrow_{35}, \rightarrow_{37}, \rightarrow_{40}, \rightarrow_{41}, \rightarrow_{42}, \rightarrow_{43}, \rightarrow_{44}, \rightarrow_{45}, \rightarrow_{47}, \rightarrow_{48}, \rightarrow_{49}, \rightarrow_{50}, \rightarrow_{52}, \rightarrow_{55}, \rightarrow_{56}, \rightarrow_{57}, \rightarrow_{58}, \rightarrow_{59}, \rightarrow_{60}, \rightarrow_{61}, \rightarrow_{62}, \rightarrow_{63}, \rightarrow_{64}, \rightarrow_{65}, \rightarrow_{66}, \rightarrow_{67}, \rightarrow_{68}, \rightarrow_{70}, \rightarrow_{71}, \rightarrow_{72}, \rightarrow_{73}, \rightarrow_{74}, \rightarrow_{76}, \rightarrow_{77}, \rightarrow_{78}, \rightarrow_{79}, \rightarrow_{80}, \rightarrow_{81}, \rightarrow_{82}, \rightarrow_{83}, \rightarrow_{84}, \rightarrow_{85}, \rightarrow_{86}, \rightarrow_{87}, \rightarrow_{88}, \rightarrow_{89}, \rightarrow_{90}, \rightarrow_{91}, \rightarrow_{92}, \rightarrow_{93}, \rightarrow_{94}, \rightarrow_{95}, \rightarrow_{96}, \rightarrow_{97}, \rightarrow_{98}, \rightarrow_{99}, \rightarrow_{102}, \rightarrow_{105}, \rightarrow_{108}, \rightarrow_{124}, \rightarrow_{125}, \rightarrow_{127}, \rightarrow_{129}, \rightarrow_{130}, \rightarrow_{132}, \rightarrow_{134}, \rightarrow_{135}, \rightarrow_{137}$  satisfy (2).

## References

- [1] Atanassov, K., *On Intuitionistic Fuzzy Sets Theory*, Springer, Berlin, 2012.
- [2] Atanassova, L., Remark on the intuitionistic fuzzy forms of two classical logic axioms. Part 1, *Annual of the "Informatics" Section, Union of Scientists in Bulgaria*, Volume 7, 2014, 24–27.
- [3] Kleene, S., *Mathematical Logic*, John Wiley & Sons, New York, 1967.