

# How to take into account monotonicity (and other properties) in centroid approach to fuzzy and intuitionistic fuzzy control

Vladik Kreinovich<sup>1</sup>  and Olga Kosheleva<sup>2</sup> 

<sup>1</sup> Department of Computer Science, University of Texas at El Paso  
500 W. University, El Paso, Texas 79968, USA  
e-mail: vladik@utep.edu

<sup>2</sup> Department of Teacher Education, University of Texas at El Paso  
500 W. University, El Paso, Texas 79968, USA  
e-mail: olgak@utep.edu

**Received:** 4 September 2025 **Accepted:** 20 November 2025 **Online First:** 27 November 2025

**Abstract:** In many application areas, there are skilled experts who excel in control and decision making. It is desirable to come up with an automated system that would use their skills to help others make similarly good decisions. Often, the experts can only formulate their skills in terms of rules that use imprecise (“fuzzy”) words from natural language like “small”. To transform these fuzzy rules into a precise control strategy, Zadeh designed special technique that he called *fuzzy*. This technique was later improved – e.g., by adding explicit information about what the experts consider *not* a good control; such addition is known as *intuitionistic fuzzy* technique. The problem that we consider in this paper is that in many cases, in addition to the fuzzy expert rules, we also have some extra knowledge about the function  $\bar{u}(x)$  – the function that describes what control to apply for a given input  $x$ . For example, it is often reasonable to require that this function is increasing: the larger  $x$ , the more control we should apply. In this paper, we use the general decision theory technique to show how this additional information can be incorporated into fuzzy and intuitionistic fuzzy control.



**Keywords:** Fuzzy sets, Intuitionistic fuzzy sets, Centroid defuzzification, Monotonicity, Decision theory.

**2020 Mathematics Subject Classification:** 03B52, 03E72.

## 1 Formulation of the problem

**Need for fuzzy control.** In many practical situations, skilled human experts can make very effective control decision, very good medical recommendations, etc. – and a few very skilled experts can make almost perfect decisions and recommendations. It is therefore desirable to design an automatic system that would incorporate the knowledge and skills of such very skilled experts – and that would help others make better decisions and recommendations.

One of the difficulties in designing such systems is that in many cases, experts cannot explain their reasoning in precise terms, they can only explain them by providing rules that are formulated in terms in imprecise (“fuzzy”) words from natural language like “small”. To transform such knowledge into precise computer-understandable form, Lotfi Zadeh came up with techniques that he called *fuzzy*; see, e.g., [3, 8, 11, 14, 15, 23].

**Fuzzy control: Main ideas.** In the original version of fuzzy techniques, each imprecise property was described by assigning, to each possible value  $x$  of the corresponding quantity, a degree  $\mu(x) \in [0, 1]$  to which the expert believes that this value satisfies the given property (e.g., that  $x$  is small). Later on, several modifications were proposed – e.g., *intuitionistic* fuzzy logic (see, e.g., [1, 22]) in which, in addition to the degree  $\mu(x)$ , the expert is asked to provide a degree  $\nu(x)$  to which, in the expert’s opinion, the degree  $x$  *does not* satisfy the given property – e.g., to what extent  $x$  is *not* small.

By using traditional fuzzy technique, for each input  $x$  and for each possible value  $u$ , we can estimate the degree  $m(x, u)$  to which, based on the expert rules,  $u$  is a reasonable control value to apply. In the intuitionistic case, this value is supplemented by the value  $n(x, u)$  to which  $u$  is *not* a reasonable control.

If all we want is to provide a recommendation to a human expert, then this information is sufficient. However, in many practical cases, we want to automate this control (or decision making). In such cases, we need to transform such fuzzy recommendations into a value  $\bar{u}(x)$  that we need to apply for the input  $x$ . Such transformation of fuzzy degrees into a single value is known as *defuzzification*. The most widely used defuzzification – known as *centroid defuzzification* – has the following form:

$$\bar{u}(x) = \frac{\int m(x, u) \cdot u \, du}{\int m(x, u) \, du}. \quad (1)$$

In the intuitionistic case, we first combine the value  $m(x, u)$  and  $n(x, u)$  into a single number – e.g., the number

$$c(x, y) = \frac{m(x, u) + (1 - n(x, u))}{2}, \quad (2)$$

and then apply centroid defuzzification to this combined membership function (see, e.g., [16]):

$$\bar{u}(x) = \frac{\int c(x, u) \cdot u \, du}{\int c(x, u) \, du}. \quad (3)$$

**Remaining question.** In many practical situations, in addition to expert rules, we have some additional information about the control function. For example, we may know that the control  $u$  should be an increasing function of  $x$ . For example, if we are describing braking, the larger the initial speed  $x$ , the more intense should be the braking.

This monotonicity is usually reflected in the corresponding rules, but – due to the fuzzy character of these rules – it is reflected only approximately. As a result, in some cases, even when rules are monotonic, the fuzzy control obtained by using the formula (1) is not everywhere monotonic; see, e.g., [21]. To avoid such situations, it is desirable to incorporate monotonicity – and other similar requirements – into fuzzy control.

**What we do in this paper.** In this paper, we show how monotonicity – and other similar properties – can be incorporated into fuzzy and intuitionistic fuzzy control.

## 2 Analysis of the problem and the resulting formulas

**Idea.** Practitioners use centroid defuzzification because it is very empirically efficient, i.e., because it is, in some reasonable sense, better than all other proposed defuzzification methods. This means that this method is optimal with respect to some reasonable criterion. So, a natural idea is to explicitly formulate this criterion – and then, solve the corresponding optimization problem under the monotonicity – (or any other) constraint.

To come up with such a criterion, let us use general decision theory.

**Decision theory: A brief reminder.** Decision theory (see, e.g., [5, 6, 9, 10, 12, 13, 17]) describes how a rational person should make a decision.

According to decision theory, a person's preferences can be described by a special function  $U(x)$  called *utility* such that out of all possible actions for which we get the result  $r_i$  with probability  $p_i$ , the decision maker selects the action for which the following expression – known as *expected utility* – attains the largest possible value:

$$U = p_1 \cdot U(r_1) + p_2 \cdot U(r_2) + \cdots + p_n \cdot U(r_n). \quad (4)$$

Utility is defined modulo adding a linear transformation – we can always choose a different starting point and a different measuring unit. To make computations simpler, let us take, as a starting point, the ideal situation when we know the exact value of the best control. In this case, in all other situations, utility will be smaller than that – i.e., its values will be negative.

**What do we need to do to apply this criterion.** This criterion is usually applied in situation when we know the probabilities and when we know the utility function. In our fuzzy control case, we do not have this information. So, to apply this approach to the fuzzy control situation, we need to estimate the probabilities and the utility. Let us do it.

**How to estimate the probabilities.** To estimate the probabilities, let us take into account that one of the natural ways to estimate the fuzzy degree is to poll several experts. For example, to decide to what extent 30° C is hot, we can ask 10 experts. If 8 of them think that 30° is hot, then we assign, as the value  $\mu(30)$  of the corresponding membership function, the degree  $8/10 = 0.8$ .

From this viewpoint, if for some input  $x$ , we have a control value  $u$  with a degree  $m(x, u)$  to which  $u$  is reasonable, we can interpret it as follows. Let us assume that we polled  $N$  experts. This means that

$$\mu(x, u) = \frac{N(x, u)}{N}, \quad (5)$$

where  $N(x, u)$  is the number of experts who believe that  $u$  is a reasonable control value for this  $x$ . Based on the formula (5), we can conclude that  $N(x, u) = N \cdot m(x, u)$ .

So, if we have values  $u_1, \dots, u_n$  with degrees  $m(x, u_1), \dots, m(x, u_n)$ , this means that:

- we have  $N \cdot m(x, u_1)$  experts who believe that  $u_1$  is a reasonable control value,
- we have  $N \cdot m(x, u_2)$  experts who believe that  $u_2$  is a reasonable control value,  $\dots$ , and
- we have  $N \cdot m(x, u_n)$  experts who believe that  $u_n$  is a reasonable control value.

So, overall, we have

$$\overline{N} \stackrel{\text{def}}{=} N \cdot m(x, u_1) + \dots + N \cdot m(x, u_n) = N \cdot (m(x, u_1) + \dots + m(x, u_n)) \quad (6)$$

opinions. We have no reasons to believe that one of these opinions is more valuable than others. So, it makes sense to assign to each of these opinions the same probability

$$p_i = \frac{1}{\overline{N}}.$$

**Comment 1.** Such an argument is often used in probability theory. It is known as *Laplace Indeterminacy Principle*; see, e.g., [7]. This principle makes perfect sense: e.g., if we have several suspects of a crime, and we have no reason to suspect one of them more than others, then it makes sense to assign, to each of them, the same *a priori* probability.

With these probabilities, the expected utility takes the following form:

$$U(u) = \frac{N \cdot m(x, u_1)}{\overline{N}} \cdot U(x, u, u_1) + \dots + \frac{N \cdot m(x, u_n)}{\overline{N}} \cdot U(x, u, u_n). \quad (7),$$

where  $U(x, u, u_i)$  is the utility of selecting the control value  $u$  when the best control value is  $u_i$ .

All the terms in this sum have the same factor  $N$  and the same denominator  $\overline{N}$  – as described by the formula (6). So, we can combine them together:

$$U(u) = \frac{N \cdot (m(x, u_1) \cdot U(x, u, u_1) + \dots + m(x, u_n) \cdot U(x, u, u_n))}{N \cdot (m(x, u_1) + \dots + m(x, u_n))}. \quad (8)$$

Dividing both numerator and denominator by  $N$ , we get a simplified expression

$$U(u) = \frac{m(x, u_1) \cdot U(x, u, u_1) + \dots + m(x, u_n) \cdot U(x, u, u_n)}{m(x, u_1) + \dots + m(x, u_n)}. \quad (9)$$

To complete this description, we need to estimate the utility values.

**How to estimate the utilities.** In most practical situations, we do not know how exactly utility depends on the control values. Such situations – when we do not know the exact form of dependence – is typical in many applications areas, e.g., in physics. In physics, the usual approach to such situations – see, e.g., [4,20] – is to take into account that the dependence is usually smooth, and usually, smooth functions can be expanded in Taylor series over some small value. To get a good approximation to the function, we can just take the sum of the first few terms in the Taylor series. In the first approximation, we usually take the smallest number of terms that are consistent with the qualitative understanding of the situations.

Let us apply this approach here. We assume that the experts are reasonably good. This means that their estimates  $u_i$  are close to the actual best value  $u$ , i.e., that the difference  $\Delta u_i \stackrel{\text{def}}{=} u_i - u$  is small. So, we can expand the expression  $U(x, u, u_i) = U(x, u, u + \Delta u_i)$  in Taylor series in terms of  $\Delta u_i$ .

We know that the utility  $U(x, u, u + \Delta u_i)$  is equal to 0 when  $u = u_i$ , i.e., when  $\Delta u_i = 0$ , and it is smaller than 0 for all other values  $\Delta u_i$ . This means that this function of  $\Delta u_i$  attains its maximum when  $\Delta u_i = 0$ . This fact implies that we cannot only keep terms which are linear in  $\Delta u_i$  – since linear functions do not attain their maxima. Thus, we need to consider quadratic terms. In general, such a quadratic expression takes the following form:

$$U(x, u, u + \Delta u_i) = a_0(x, u) + a_1(x, u) \cdot \Delta u_i + a_2(x, u) \cdot (\Delta u_i)^2. \quad (10)$$

The fact that this value is equal to 0 when  $\Delta u_i = 0$  implies that  $a_0(x, u) = 0$ . The fact that the function (10) attains its maximum for  $\Delta u_i = 0$  implies that the derivative of the expression (10) is equal to 0 when  $\Delta u_i = 0$  – which implies that  $a_1(x, u) = 0$ . Thus:

$$U(x, u, u + \Delta u_i) = a_2(x, u) \cdot (\Delta u_i)^2. \quad (11)$$

The fact that all utility values are non-negative means that  $a_2(x, u) < 0$ .

We can simplify the expression (11) even further if we take into account that, as we have mentioned earlier, the utility usually smoothly depends on  $u$  – which means that the function  $a_2(x, u)$  also smoothly depends on  $u$ . So, we can select some value  $u_0$  close to all the expert estimates, so that  $a(x, u) = a(x, u + \Delta u)$ , where we denoted  $\Delta u \stackrel{\text{def}}{=} u - u_0$ . By applying the same Taylor approximation idea as before, we conclude that in the first approximation, we have

$$a_2(x, u) = a_2(x, u_0 + \Delta u) = a_2(x, u_0) + a_3(x, u_0) \cdot \Delta u. \quad (12)$$

Substituting the expression (12) into the formula (11), we get

$$U(x, u, u + \Delta u_i) = a_2(x, u_0) \cdot (\Delta u_i)^2 + a_3(x, u_0) \cdot \Delta u \cdot (\Delta u_i)^2. \quad (13)$$

The last term in this expression is already cubic in terms of small differences  $\Delta u$  and  $\Delta u_i$ . Since we are only considering quadratic approximation, we can safely ignore this terms and get:

$$U(x, u, u_i) = U(x, u, u + \Delta u_i) = a_2(x, u_0) \cdot (\Delta u_i)^2 = a_2(x, u_0) \cdot (u_i - u)^2. \quad (14)$$

**Now, we get the final expression for utility.** Substituting the expression (14) into the formula (9), we conclude that

$$U(u) = \frac{m(x, u_1) \cdot a_2(x, u_0) \cdot (u_1 - u)^2 + \cdots + m(x, u_n) \cdot a_2(x, u_0) \cdot (u_n - u)^2}{m(x, u_1) + \cdots + m(x, u_n)}. \quad (15)$$

By definition of utility, for each  $x$ , we need to select the control value  $u$  for which the utility is the largest possible. If we multiply all the values of the objective function by a positive constant, this does not change which value is larger and which is smaller: e.g., who is the richest person does not change whether we count his/her amount in US dollars or in Euros. Thus, for the purpose of maximizing utility, we can multiply all the values of the utility function by the denominator of the expression (15) and divide by the absolute value  $|a_2(x, u_0)|$ . This way, we get the equivalent maximizing function

$$\overline{U}(u) \stackrel{\text{def}}{=} -(m(x, u_1) \cdot (u_1 - u)^2 + \cdots + m(x, u_n) \cdot (u_n - u)^2). \quad (16)$$

Maximizing this expression is equivalent to minimizing its opposite

$$-\overline{U}(u) = m(x, u_1) \cdot (u_1 - u)^2 + \cdots + m(x, u_n) \cdot (u_n - u)^2. \quad (17)$$

In the continuous case, the sum becomes the integral, so we want to minimize the expression

$$\int m(x, u) \cdot (u - \bar{u})^2 du. \quad (18)$$

**Comment 2.** From the computational viewpoint, the minimized expressions (17) and (18) are a particular case of the general Least Squares problem (see, e.g., [19]), when we minimize the sum of squares of some expressions.

**If there are no additional constraints, this expression leads exactly to centroid defuzzification.** Indeed, if we differentiate the expression (17) with respect to  $u$  and equate the derivative to 0, we conclude that

$$2m(x, u_1) \cdot (u - u_1) + \cdots + 2m(x, u_n) \cdot (u - u_n) = 0. \quad (19)$$

If we divide both sides by 2, move all the terms not containing  $u$  to the right-hand side, and group together all the terms proportional to  $u$  in the left-hand side, then we get the following linear equation with one unknown:

$$(m(x, u_1) + \cdots + m(x, u_n)) \cdot u = m(x, u_1) \cdot u_1 + \cdots + m(x, u_n) \cdot u_n. \quad (20)$$

Thus:

$$u = \frac{m(x, u_1) \cdot u_1 + \cdots + m(x, u_n) \cdot u_n}{m(x, u_1) + \cdots + m(x, u_n)}. \quad (21)$$

In the continuous case, when sums turn into integrals, we get exactly the centroid defuzzification (1).

**What to do if we have an additional constraint on the dependence  $\bar{u}(x)$  – e.g., monotonicity.** In the previous part of the section, we only considered utility corresponding to a single input  $x$ . If we impose an additional restriction that relates controls corresponding to different inputs, then

we need to consider the overall utility, i.e., the sum – or integral – of the expression (15) over all  $x$ .

Similarly to what we argued about the possibility to ignore, in this approximation, the dependence of  $a_2(x, u)$  on  $u$ , we can argue that – at least for some range of inputs – we can also ignore the dependence on  $x$ , i.e., replace  $a_2(x, u_0)$  with  $a_2(x_0, u_0)$ . In this case, maximizing the overall utility for all possible values of  $x$  is equivalent to minimizing the following integral

$$\int_x \int_u m(x, u) \cdot (u - \bar{u}(x))^2 du dx. \quad (22)$$

So, if we know some additional constraint(s) on the function  $\bar{u}(x)$ , e.g., that this dependence should be monotonic, then we should find, among all the functions  $\bar{u}(x)$  that satisfy these constraints, the one for which the expression (22) attains the smallest possible value.

**How to actually find such control: case of monotonicity.** The minimized function (22) is still the sum (integral) of squares, so it is still a particular case of the general Least Squares problem. For the case when the restriction is that the function  $\bar{u}(x)$  is a (non-strictly) increasing function of  $x$ , the problem becomes: to find an increasing solution to the Least Squares problem. In data processing, this problem is known as *isotonic regression*. There exist efficient algorithms for solving this problem; see, e.g., [2, 18].

**What to do in the intuitionistic fuzzy case.** In the intuitionistic fuzzy case, for each  $x$  and  $u$ , in addition to

$$N \cdot m(x, u)$$

experts who believe that  $u$  is a reasonable control value, there are also  $N \cdot n(x, u)$  experts who believe that  $u$  is *not* a reasonable control value.

To take their opinions into account, we need to describe it in terms of utility, i.e., we need to describe the utility  $V(x, u, u_i)$  corresponding to the case when the actual best control value is  $u$  but the  $i$ -th expert says that  $u_i$  is not a good value. Similarly to the fuzzy case, we can select a starting point for the utility as the point for which  $u_i = u$ . The difference is that now this is the worst case, when the expert is completely wrong. So, in this case, for all  $u_i \neq u$ , the utility is larger. Thus, the function  $V(x, u, u_i)$  should now attain its *minimum* for  $u_i = u$ . Similarly to the fuzzy case, this means that we cannot restrict ourselves to linear terms in the Taylor expansion, we need to take quadratic terms into account. If we restrict ourselves to quadratic terms, then, similarly to the fuzzy case, we get

$$V(x, u, u_i) = b_2(x, u_0) \cdot (u - u_i)^2, \quad (23)$$

where this time, the expression  $b_2(x, u_0)$  is positive.

If we now add up all terms corresponding to positive and negative opinions and perform the same simplifications as in the fuzzy case, we conclude that maximizing utility is equivalent to minimizing the expression

$$m(x, u_1) \cdot (u - u_1)^2 - r \cdot n(x, u_1) \cdot (u - u_1)^2 + \dots + m(x, u_n) \cdot (u - u_n)^2 - r \cdot n(x, u_n) \cdot (u - u_n)^2, \quad (24)$$

where  $r \stackrel{\text{def}}{=} b(x, u_0)/|a(x, u_0)|$ . In other words, we want to minimize the expression

$$c(x, u_1) \cdot (u - u_1)^2 + \cdots + c(x, u_n) \cdot (u - u_n)^2, \quad (25)$$

where  $c(x, u_i) = m(x, u_i) - r \cdot n(x, u_i)$  is a combination of positive and negative membership functions. In the continuous case, this means minimizing the expression

$$\int c(x, u) \cdot (u - \bar{u})^2 du. \quad (26)$$

Similarly to the fuzzy case, in the absence of additional constraints, minimizing the expressions (25)–(26) is equivalent to applying centroid defuzzification to the combined membership function. In the case of constraints, we need to find the function  $\bar{u}(x)$  that, among all the functions that satisfy the given constraints, minimizes the expression

$$\int_x \int_u c(x, u) \cdot (u - \bar{u})^2 du dx. \quad (27)$$

For the case of monotonicity, we can similarly use isotonic regression techniques from [2, 18] to find the corresponding function.

### 3 Conclusion

Let us summarize our results. In this paper, we discussed a problem of how to take into account additional a priori constraints on the control function  $\bar{u}(x)$  in fuzzy and intuitionistic fuzzy control. To solve this problem, we used the general techniques of decision theory. By applying these techniques, we came up with the following conclusions.

In the fuzzy case, when we have a membership function  $m(x, u)$  – that describes to what extent  $u$  is an appropriate control for the input  $x$  – to find the corresponding control, we need to find, among all the functions  $\bar{u}(x)$  that satisfy the given constraints, the function that minimizes the following expression:

$$\int_x \int_u m(x, u) \cdot (u - \bar{u}(x))^2 du dx. \quad (22)$$

In the intuitionistic fuzzy case, when we also have a negative membership function  $n(x, u)$  that describes to what extent  $u$  is *not* an appropriate control for the input  $x$ , we need to minimize the following expression:

$$\int_x \int_u c(x, u) \cdot (u - \bar{u}(x))^2 du dx, \quad (27)$$

where  $c(x, u) = m(x, u) - r \cdot n(x, u)$  is a combination of positive and negative membership functions.

In the important case when the constraint means that the function  $\bar{u}(x)$  should be an increasing function of  $x$ , we can use efficient algorithms of isotonic regression [2, 18] to minimize the above expressions.



## Acknowledgements

This work was supported in part by the National Science Foundation grants 1623190 (A Model of Change for Preparing a New Generation for Professional Practice in Computer Science), HRD-1834620 and HRD-2034030 (CAHSI Includes), EAR-2225395 (Center for Collective Impact in Earthquake Science C-CIES), and by the AT&T Fellowship in Information Technology.

It was also supported by a grant from the Hungarian National Research, Development and Innovation Office (NRDI), by the Institute for Risk and Reliability, Leibniz Universitaet Hannover, Germany, and by the European Union under the project ROBOPROX (No. CZ.02.01.01/00/22 008/0004590).

The authors are greatly thankful to Dr. Krassimir Atanassov for his encouragement.

## References

- [1] Atanassov, K. (1999). *Intuitionistic Fuzzy Sets: Theory and Applications*. Springer-Verlag, Heidelberg.
- [2] Barlow, R. E., Bartholomew, D. J., Bremner, J. M., & Brunk, H. D. (1972). *Statistical Inference under Order Restrictions. The Theory and Application of Isotonic Regression*. Wiley, New York.
- [3] Belohlavek, R., Dauben, J. W., & Klir, G. J. (2017). *Fuzzy Logic and Mathematics: A Historical Perspective*. Oxford University Press, New York.
- [4] Feynman, R., Leighton, R., & Sands, M. (2005). *The Feynman Lectures on Physics*. Addison Wesley, Boston, Massachusetts.
- [5] Fishburn, P. C. (1969). *Utility Theory for Decision Making*. John Wiley & Sons Inc., New York.
- [6] Fishburn, P. C. (1988). *Nonlinear Preference and Utility Theory*. The John Hopkins Press, Baltimore, Maryland.
- [7] Jaynes, E. T., & Bretthorst, G. L. (2003). *Probability Theory: The Logic of Science*. Cambridge University Press, Cambridge, UK.
- [8] Klir, G., & Yuan, B. (1995). *Fuzzy Sets and Fuzzy Logic*. Prentice Hall, Upper Saddle River, New Jersey.
- [9] Kreinovich, V. (2014). Decision making under interval uncertainty (and beyond), In: Guo, P., & Pedrycz, W. (Eds.). *Human-Centric Decision-Making Models for Social Sciences*. Springer Verlag, pp. 163–193.
- [10] Luce, R. D., & Raiffa, R. (1989). *Games and Decisions: Introduction and Critical Survey*. Dover, New York.

- [11] Mendel, J. M. (2024). *Uncertain Rule-Based Fuzzy Systems: Introduction and New Directions*. Springer, Cham, Switzerland.
- [12] Nguyen, H. T., Kosheleva, O., & Kreinovich, V. (2009). Decision making beyond Arrow's 'impossibility theorem', with the analysis of effects of collusion and mutual attraction. *International Journal of Intelligent Systems*, 24(1), 27–47.
- [13] Nguyen, H. T., Kreinovich, V., Wu, B., & Xiang, G. (2012). *Computing Statistics under Interval and Fuzzy Uncertainty*. Springer Verlag, Berlin, Heidelberg.
- [14] Nguyen, N. T., & Walker, E. A. (2019). *A First Course in Fuzzy Logic*. Chapman and Hall/CRC, Boca Raton, Florida.
- [15] Novák, V., Perfilieva, I., & Močkoř, J. (1999). *Mathematical Principles of Fuzzy Logic*. Kluwer, Boston, Dordrecht.
- [16] Radhika, C., & Parvathi, R. (2016). Defuzzification of intuitionistic fuzzy sets. *Notes on Intuitionistic Fuzzy Sets*, 22(5), 19–26.
- [17] Raiffa, R. (1997). *Decision Analysis*. McGraw-Hill, Columbus, Ohio.
- [18] Robertson, T., Wright, F. T., & Dykstra, R. L. (1988). *Order Restricted Statistical Inference*. Wiley, New York.
- [19] Sheskin, D. J. (2011). *Handbook of Parametric and Nonparametric Statistical Procedures*. Chapman and Hall/CRC, Boca Raton, Florida.
- [20] Thorne, K. S. & Blandford, R. D. (2021). *Modern Classical Physics: Optics, Fluids, Plasmas, Elasticity, Relativity, and Statistical Physics*. Princeton University Press, Princeton, New Jersey.
- [21] Van Broekhoven, E., & De Baets, B. (2005). A linguistic fuzzy model with a monotone rule base is not always monotone. *Proceedings of the EUSFLAT-LFA'05*, Barcelona, Spain, 2005, pp. 530–535.
- [22] Vassilev, P. M., & Atanasov, K. T. (2019). *Modifications and Extensions of Intuitionistic Fuzzy Sets*. "Prof. Marin Drinov" Academic Publishing House, Sofia.
- [23] Zadeh, L. A. (1965). Fuzzy sets. *Information and Control*, 8, 338–353.